95.02.03

INSECT-KURE (R)

INSECTICIDE

LIQUID

FOR BARN AND LIVESTOCK

READY TO USE

WILL CONTROL: Flies, Mosquitoes, Gnats

AGRICULTURAL

SHAKE WELL BEFORE USING

READ THE LABEL BEFORE USING

10 LITRES

GUARANTEE:

REGISTRATION NUMBER 23820

PEST CONTROL PRODUCTS ACT

Exclusive Distributor

Aerokure International Inc. Box 22, Sherbrooke (Quebec) Canada JlH 5H5

DIRECTIONS

Space Spray: To control house fly, stable fly, mosquitoes and gnats, in hog, dairy and beef barns, milking parlors and milk

rooms. Close doors and windows. Apply as a fog or fine mist directing spray upwards to all parts of the room especially windows and other light sources, using U.L.V. equipment which produces a find mist such as Cyclone. Use 200-300 mL of solution/100 m3. Leave the room closed for 20 minutes before ventilation. The ideal time of the day for spraying is early morning. Spray only when temperature is under 32 deg. C. This product is non-corrosive. It must be protected from sun rays. Do not forget to close container after use. Repeat application as necessary. Remove all animals from barn before using as a space spray. Ventilate treated areas thoroughly for at least 30 minutes before animal re-entry.

LIVESTOCK SPRAY

To control horse fly, house fly, mosquitoes, stable fly, horn fly, face fly, black fly, lice, flea, ticks, on beef cattle, dairy cattle (lactating, non lactating), hogs and horses. Spray animal, not more than once daily, using a sprayer which produces a fine mist such as a cold aerosol fogger. Apply at a rate of 30-60 mL/animal. Spray should be enough to wet ends of the hair only. Do not wet the hair on skin. Take care not to spray in animal's eyes. Treat dairy cattle before milking but do not spray directly on the udders. Wash udders before milking. After one week, treatments may be reduced to once every second day.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN

Apply only when dairy barn is not in operation. Remove all food before treatment. Avoid inhalation of spray mist. Do not contaminate feed or water troughs. Do not remain in treated areas. Ensure proper ventilation after sealed areas have been treated. Cover milk and milking utensils to prevent contamination from spray and dead insects. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before re-use. Avoid contact with skin and eyes. Wash after using.

DISPOSAL

- Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
- Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- Make the empty container unsuitable for further use.
- Dispose of the container in accordance with provincial requirements.
- For information on the disposal of unused, unwanted product

and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

FIRST AID

If in eyes, or on skin, flush with clean water. If irritation persists, obtain medical attention. If accidentally swallowed, obtain medical attention at once.

NOTICE TO USER

This product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))