

96/2001

WEEDAWAY (R)

2,4-DB 400

10 LITRES

EMULSIFIABLE CONCENTRATE HERBICIDE

AGRICULTURAL

FOR WEED CONTROL IN SEEDLING ALFALFA, BIRDSFOOT TREFOIL AND
PASTURES

READ THE LABEL BEFORE USING

GUARANTEE: 2,4-DB 400 g/L (Present as Ester)

REGISTRATION NO: 21929
PEST CONTROL PRODUCTS ACT

WARNING POISON

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9

(R) T.M. Reg'd.

LABEL INDEX 2,4-DB 400	
Seedling Alfalfa and Birdsfoot Trefoil (direct or underseeded) GO TO SECTION 1	Weeds Controlled and Rate GO TO SECTION 6
Seedling Clover (direct or underseeded) GO TO SECTION 2	Tank Mix with MCPA (To improve wild mustard control in seedling alfalfa) GO TO SECTION 7
Cereals(Wheat, Oats, Barley) GO TO SECTION 3	Filling instructions GO TO SECTION 8
Field Corn GO TO SECTION 4	Precautions GO TO SECTION 9

Pastures Containing Forage Legumes GO TO SECTION 5	First Aid	GO TO SECTION 10
	Rinsing and Disposal	GO TO SECTION 11

SECTION 1	
Crop:	SEEDLING ALFALFA AND BIRDSFOOT TREFOIL (direct or underseeded) NOTE: Do Not Apply if Crop is to be used for Seed Production Later in the Season.
Treatment Stage:	Treat from 1st to 4th trifoliate leaf stage.
GO TO SECTION 6	

SECTION 2	
Crop:	SEEDLING CLOVERS (direct or underseeded) NOTE: Do Not Spray Sweet Clover. Red and Alsike Clovers may be damaged.
Treatment Stage:	Treat as soon as possible after the first trifoliate leaf appears.
GO TO SECTION 6	

SECTION 3	
Crop:	CEREALS (WHEAT, OATS, BARLEY)
Treatment Stage:	Apply after the crop has 5 fully expanded leaves but before the early flag leaf stage. Oats will be damaged if treated before 5 leaf stage.
GO TO SECTION 6	

SECTION 4	
Crop:	FIELD CORN
Treatment Stage:	Apply after crop is at least 40 cm high but before tasselling. Use drop nozzles to avoid spraying corn leaves.
GO TO SECTION 6	

SECTION 5	
Crop:	PASTURES CONTAINING FORAGE LEGUMES
Treatment Stage:	Treat after cutting or grazing when growth is not above 7.5 cm in height. Damage may occur if crop is sprayed beyond this stage.
GO TO SECTION 6	

SECTION 6		
Annual Weeds Controlled:		Rate:
Wormseed Mustard, Wild Mustard, Ball Mustard, Stinkweed, Ragweed, Redroot Pigweed, Shepherd's Purse, Lamb's-quarters.		2.75 L/ha
Oakleaved Goose Foot, Wild Buckwheat, Wild Radish (NOTE: Best control occurs when weeds sprayed in the 1-2 leaf stage.)		3.5 L/ha
Lady's-thumb, Green Smartweed. Seedlings stunted only, but competitive ability is reduced.		4.25 L/ha
Perennial Weeds:	Spray Stage:	Rate:

Bull Thistle	Rosette to early bud)	2.75-4.25 L/ha
Perennial Sow Thistle	Rosette)	
Canada Thistle	Top growth only. Spray) from 15 cm to early bud)	
Curled Dock	Early growth)	
Plantain	Before flowering)	
Field Bindweed	Top growth reduced. For) best results, spray in) late Summer)	
Yellow Rocket	Spray between 3rd week in September and middle of October)	
Chicory	Rosette stage)	
Narrow Leafed Hawks-beard	Rosette stage in fall after legume growth ends	3.5 L/ha
Dandelion	Top growth only. Spray before bud stage.	4.25 L/ha
Horsetail	Top growth only. Spray when 10-12 cm high.	4.25 L/ha
<p>IMPORTANT NOTICE:</p> <p>Use lower rate on susceptible weeds in the seedling stage. Use the higher rates for older, heavy weed infestations and under poor growing conditions.</p> <p>Do not exceed recommended rate for each crop. Apply product in warm weather.</p> <p>Do not apply under drought conditions.</p>		
GO TO SECTION 8		

SECTION 7	
Tank Mix with MCPA:	To improve wild mustard and narrow leafed hawksbeard control in seeding Alfalfa.

Treatment Stage:	Treat mustard beyond four leaf stage. Apply 2,4-DB 400 alone on mustard at 4 leaf stage or less.	
Rate:	2,4-DB 400 + MCPA Amine 500	2 L/ha + 70 mL/ha
GO TO SECTION 8		

SECTION 8

FILLING SPRAY TANK

1. Fill tank 1/2 full with water.
2. Add 2,4-DB 400
3. When MCPA Amine 500 is used, add and mix before adding 2,4-DB 400.
4. Complete filling of tank with water.
5. Use 150-200 lites of water per hectare.
6. Do not exceed 275 kPa pressure.

GO TO SECTION 9

SECTION 9

Precautions:

1. Keep out of reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapors. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally hot days over 27 deg. C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application exercise special care:
 - 1) Do not spray in winds exceeding 8 km per hour.
 - 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
 - 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.

9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. DO not re-use empty containers.

GO TO SECTION 10

SECTION 10

First Aid:

1. In case of poisoning, call a physician immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Obtain medical attention.
4. If swallowed, do not induce vomiting, but rush patient to nearest hospital or doctor's office. Take the pesticide label with you.

GO TO SECTION 11

SECTION 11

Disposal Information:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

Notice to User: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

Notice to Buyer: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of handling of this product and accepts the product on that condition.

96/2001

[[LABEL]]

WEEDAWAY (R)

2,4-DB 400

10 LITRES

EMULSIFIABLE CONCENTRATE HERBICIDE

AGRICULTURAL

FOR WEED CONTROL IN SEEDLING ALFALFA, BIRDSFOOT TREFOIL AND
PASTURES

READ THE LABEL BEFORE USING

GUARANTEE: 2,4-DB 400 g/L (Present as Ester)

REGISTRATION NO: 21929
PEST CONTROL PRODUCTS ACT

WARNING POISON

Interprovincial Cooperative Limited,
P.O. Box 1050, Saskatoon, Saskatchewan S7K 3M9

(R) T.M. Reg'd.

PRECAUTIONS:

1. Keep out of reach of children.
2. Avoid contact with eyes, skin and clothing. May cause skin irritation.
3. Do not breathe spray mist or vapors. Avoid spray or vapor drifts onto susceptible plants such as grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, turnips, cauliflower, cabbage, alfalfa, clover, peas, potatoes, ornamental trees, shrubs or flowers, fruit trees, lawns, most vegetable crops or other desirable vegetation.
4. Do not spray when there is any danger of wind or on exceptionally hot days over 27 deg. C.
5. Drift from spray may be reduced by using high volume sprays under low pressure, coarse sprays, and drop nozzles. Use special care in airplane application where danger from drift is greater.
6. For aerial application exercise special care:

- 1) Do not spray in winds exceeding 8 km per hour.
- 2) Do not spray in dead calm near sensitive plants. A cloud of suspended droplets may drift onto sensitive plants when the wind comes up.
- 3) Spray only when the wind is blowing away from a sensitive crop, shelter belt or garden.
7. Do not contaminate any body of water. Avoid contamination of foods.
8. Do not use in a greenhouse.
9. Keep in original container during storage.
10. Do not store near or in same room as seeds, feeds, fertilizers or pesticides.
11. Keep away from fire or open flame and other sources of heat.
12. Clean spray equipment thoroughly after use. Do not use spray equipment for other purposes. DO not re-use empty containers.

DISPOSAL INFORMATION:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

FIRST AID:

1. In case of poisoning, call a physician immediately.
2. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water.
3. If in eyes, flush with water immediately and continue for 5-10 minutes. Get medical attention.
4. If swallowed, do not induce vomiting. Take patient to nearest hospital or doctor's office. Take the pesticide label with you.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))