

Pouch, Main panel

NET CONTENTS: 100 g

RUSE II

RODENTICIDE
PELLETS PLACE PACKS

Kill Rats and Mice with a Single Feeding
Kills Warfarin-Resistant Norway Rats

DOMESTIC

GUARANTEE: BROMADIOLONE 0.005%

Warning, contains the allergens peanut and wheat.

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL AND OUTER CONTAINER BEFORE USING

REGISTRATION NO.25772 - PEST CONTROL PRODUCTS ACT

WARNING



POISON

Vétoquinol N.-A. Inc.
2000, ch. Georges, Lavaltrie
QC, Canada J5T 3S5
1-800-363-1700

Pouch, Secondary panel

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. This material may reduce the clotting ability of the blood and cause bleeding. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID AND INSTRUCTIONS: If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object, repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. For all cases of human ingestion immediately notify a physician or poison control center. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

DISPOSAL: Do not reuse empty container. Wrap and dispose of empty container with household garbage.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

Box, Main panel

NET CONTENTS: 1 kg (10 X 100 g)

RUSE II

RODENTICIDE
PELLETS PLACE PACKS

Kill Rats and Mice With a Single Feeding
Kills Warfarin-Resistant Norway Rats

DOMESTIC

GUARANTEE: BROMADIOLONE 0.005%

Warning, contains the allergens peanut and wheat.

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING

REGISTRATION NO.25772 - PEST CONTROL PRODUCTS ACT

WARNING



POISON

Vétoquinol N.-A. Inc.
2000, ch. Georges, Lavaltrie
QC, Canada J5T 3S5
1-800-363-1700

Box, Secondary panel

DIRECTIONS FOR USE: Ruse II Pellets is for the control of Norway rats, roof rats and house mice indoors and against the outside walls of buildings.

USE LIMITATIONS: Baits **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Place bait in locations where rats and mice travel and will most likely find and consume the bait. Placement along walls, in corners, by gnawed openings, in and around burrows, and in concealed locations where rats and mice have been observed. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

APPLICATION DIRECTIONS: For Norway Rats (*Rattus norvegicus*) and roof rats (*Rattus rattus*): Place 1 to 4 packs of bait (usually at intervals of 5 to 10 m) within the treatment area. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. For House Mice (*Mus musculus*): Place 30 g of bait at intervals of 2 to 4 m within the treatment area. Sixty (60) g of bait per placement may be necessary at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS & MICE: Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations as needed.

PRECAUTIONS: **KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.** May be harmful or fatal if swallowed or absorbed through the skin. This material may reduce the clotting ability of the blood and cause bleeding. Do not open pre-measured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.** This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID: If in eyes, flush with water for 15 minutes. If on skin, wash affected area thoroughly with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of throat with a finger or blunt object, repeat until vomitus is clear. Do not induce vomiting or give anything by mouth to an unconscious person. For all cases of human ingestion immediately notify a physician or poison control center. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE AND DISPOSAL: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals. Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and dispose of empty container with household garbage.

RODENT DISPOSAL: Dispose of dead rodents in garbage or by burying.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.