SANEX

MOXY

GARDEN SPRAY CONCENTRATE

Insecticide concentrate containing Methoxychlor 25% to control Fruit, Vegetable and Ornamental insects, Mosquitoes and Blackflies

DOMESTIC

DANGER

POISON

Read label before using

500 mL

GUARANTEE:
Methoxychlor 240 g/L

REG. NO. 12733 P.C.P. ACT

Manufactured by

SANEX INC.

2695 Slough Street, Mississauga, Ontario L4T 1G2

Branches Throughout North America

DIRECTIONS FOR USE

Roses, flowers, shrubs, ornamentals, shade trees: Controls Beetles, Caterpillars, White Flies, Thrips, Coopers, Weevils, Rose Chafer, Box Elder Bug and other leaf-eating insects. Use 40 mL/L water and spray thoroughly. Avoid heavy application to elms and evergreens.

Apple, pear, cherry, peach, prune, or plum: Use 6-12.5 mL/L water and spray thoroughly. Controls Plum Curcurlio and Cherry Fruit Fly. Apply at petal fall and repeat at 7-14 day intervals as

necessary. Do not apply to apples, pears, and cherries within 7 days of harvest or to peaches within 21 days of harvest.

Strawberries, gooseberries, currants: For Strawberry Weevil and Currant Fruit Fly, apply 6-12.5 mL as a thorough cover spray when insects appear. Do not apply within 14 days of harvest.

Vegetables: Use 6-12.5 mL/L water and spray for thorough coverage. Repeat in 7 to 10 days as required.

Snap Beans: Controls Mexican Bean Beetle, Leafhoppers.

Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Carrots, Lettuce, Radishes, Spinach, Turnips, Pepper, Eggplant, Cucumber: Controls Flea Beetles, Leafhoppers, Blister Beetles, Cucumber Beetles. Do not apply within 1 day of harvest on carrots, Cucumber, Eggplant, Kohlrabi Peppers, Turnips: 3 days of harvest on Cabbage; 7 days on Beans, Beets, Cauliflower and Radishes; 14 days on Broccoli, Brussels Sprouts, Collards, Lettuce and Spinach.

Potatoes: Controls Colorado Potato Beetle, Flea Beetle, Leafhoppers, Blister Beetle. Do not apply within 7 days of harvest.

Tomatoes: Controls mosquitoes, Gnats, Wasps, Hornets, Flying Moths, Sandflies, Silverfish, Crickets, Clover Mites, Carpet Beetles, Stored Product Insects. Mix 25 mL/L of water and use 200-400 mL per 100 m3 of space. Use as a contact spray for crawling insects. Avoid contamination of food utensils and surrounding surfaces that may contact food. Remove pets from area before spraying.

Adult Mosquitoes & blackflies around buildings & cottage sites: Apply 12.5 mL per litre of water and spray to cover 100 squire meters of ground area thoroughly using hand or power sprayer. Repeat applications may be necessary. Blackfly control may not be as good as mosquito control. Do not spray marshes, swamps, and open water.

PRECAUTION: Avoid contact with skin. Avoid breathing mist. Avoid contamination of feed and foodstuffs. Keep spray away from fishbearing waters and bird feeding stations. Do not store near heat or open flame.

FIRST AID: For external contact, wash with soap and water. If in eyes, flush with plenty of water. If swallowed, cause vomiting by placing a finger at the back of the throat. Repeat until vomit fluid is clear. Obtain prompt medical attention.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arisings from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))