

12-JUL-1996

ISK BIOSCIENCES(TM)

TERSAN(R) 1991

TURF FUNGICIDE

WETTABLE POWDER FOR TURF

DOLLAR SPOT
BROWN PATCH
PINK SNOW MOLD

COMMERCIAL

READ THE LABEL BEFORE USING

GUARANTEE:
Benomyl 50%

REGISTRATION NO: 11061.01
PEST CONTROL PRODUCTS ACT

Net Contents: 2 kg

PRECAUTION!

May irritate eyes, nose, throat, and skin.

KEEP OUT OF REACH OF CHILDREN.

Avoid breathing dust or spray mist.
Avoid contact with skin, eyes and clothing.
Keep away from fire or sparks.

FIRST AID: In case of contact, flush skin or eyes with plenty of water; for eyes, get medical attention or contact poison control centre.

For medical emergencies involving this product, call
toll free 1-800-441-3637.

ISK Biosciences Corporation
5966 Heisley Road
P.O. Box 8000
Mentor, Ohio 44061-8000

DIRECTIONS

"Tersan" 1991 Turf Fungicide should be used only in accordance with recommendations on this label.

"Tersan" 1991 Turf Fungicide is recommended for use as a spray for the control of Dollar Spot, Brown Patch, Fusarium Blight and Pink Snow Mold. "Tersan" 1991 provides both curative (eradicant) action and residual protective action.

Apply recommended amounts in sufficient water to obtain thorough coverage, usually 10-22 L/100 sq. m. When treating golf greens, always treat aprons and approaches.

TURF DISEASES

Dollar Spot (*Sclerotinia homoeocarpa*)--Apply "Tersan" 1991 at rate of 30 g/100 m² at earliest appearance of disease. Repeat at 2 week intervals or when disease begins to reappear.

Brown Patch (*Rhizoctonia solani*)--Apply "Tersan" 1991 at rate of 60 g per 100 m² at earliest appearance of disease and continue at 10 to 14 day intervals as long as needed. When conditions are unusually favourable for development of disease, reduce interval to 5 to 7 days.

Pink Snow Mold (*Fusarium nivale*)

B.C. AND EASTERN CANADA

Fall Protection: Apply 60 g per 100 m² at first sign of disease. Repeat two weeks later if disease persists.

Over Winter Protection: Apply 125 g per 100 m² during November.

A second application in early spring at 60 g may be needed if infection re-appears.

NOTE: Where Melting Out, (Leaf Spot, Blight) (*Helminthosporium* spp.) are a problem, apply "Tersan" LSR Turf Fungicide two or three times in the early spring at rate of 90-125 g/100 m². at 7 to 10 day intervals.

IMPORTANT: Never allow "Tersan" 1991 to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of "Tersan" 1991 as a fungicide. Keep container closed when not in use. Do not re-use container.

Do not graze or feed clippings from treated areas to livestock.

Do not contaminate water or spray in areas where surface flow may enter water courses.

STORAGE AND DISPOSAL

STORAGE:

Store product in original container only, away from fertilizer, food, or feed. Not for use or storage in or around the home. Keep container closed.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

Purchase of this material does not confer any rights under patents of countries outside of Canada.

(R)Tersan is a registered trademark of E.I. du Pont de Nemours and Company

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))