

96.10.16

((POUCH))

SANEX(R)
LIQUID BROMONE(R)
COMMERCIAL
RODENTICIDE BAIT
NET CONTENTS: 50 mL

GUARANTEE:

Bromadiolone 0.011%

READ THE LABEL AND OUTER CARTON BEFORE USING

REGISTRATION NO. 21179
PEST CONTROL PRODUCTS ACT

WARNING POISON

MIX POUCH CONTENTS WITH 450 ML OF WATER

SANEX AGRO INC.
Dundas, Ontario
L9H 3H3

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, DOMESTIC ANIMALS, PETS AND WILDLIFE. Before using, see outer carton for full PRECAUTIONS regarding placement. storage and hygiene.

FIRST AID

This product contains an anti-coagulant chemical which, if taken accidentally by humans, domestic animals, pets or wildlife, may reduce the clotting ability of the blood and haemorrhage may result. If in eyes, flush with plenty of water for 15 minutes. If on skin, wash thoroughly with soap and water. If swallowed, give

patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object. Repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

DISPOSAL

Dispose of the container in accordance with provincial requirements.

((CARTON))

SANEX(R)

LIQUID BROMONE(R)

COMMERCIAL

RODENTICIDE BAIT

For control of rats and mice
One feeding kills rats and mice
Kills warfarin resistant Norway rats

NET CONTENTS: 12 x 50 mL

GUARANTEE:

Bromadiolone 0.011%

READ THE LABEL BEFORE USING

REGISTRATION NO. 21179
PEST CONTROL PRODUCTS ACT

WARNING POISON

MIX POUCH CONTENTS WITH 450 ML OF WATER

SANEX AGRO INC.
36 Head Street
Dundas, Ontario
L9H 3H3

REFER TO AND RETAIN THIS CARTON FOR COMPLETE USE AND HANDLING INSTRUCTIONS.

For control of Norway rats (*Rattus norvegicus*) Roof rats (*Rattus rattus*) and house mice (*Mus musculus*) in and around homes, industrial, commercial, agricultural and public buildings.

Selection of treatment areas:

Determine areas where rats and/or mice will most likely find and drink the bait. These areas include along the walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls or in locations where rodents and their signs have been observed. Remove as much food as possible from treatment area.

MIXING DIRECTIONS:

Mix complete contents of each 50 mL pouch with 450 mL of tap water.

APPLICATION DIRECTIONS:

Norway and Roof rats:

Place a minimum of 500 mL of Sanex Liquid Bromone mixture (50 mL Liquid Bromone plus 450 mL water) in a dispenser such as a chick fountain or equivalent device at each placement. Placements should be at intervals of 5-8 metres. Maintain an uninterrupted supply of fresh liquid bait for at least 10 days.

House Mice:

Place diluted Sanex Liquid Bromone bait in dispensers such as chick fountains or equivalent and position the dispensers at intervals of 2-3 metres. Maintain an uninterrupted supply of fresh liquid bait for at least 15 days.

General

Replace contaminated or spoiled bait immediately or every 4-5 days. Collect and dispose of all dead rodents and unconsumed bait properly. Repeat treatment when infestation occurs. Resistant infestations may require establishing permanent bait stations.

For areas not directly related to food processing: Use only in non-food or non-feed areas where feed, food packaging and handling equipment are never opened or exposed. For areas where

feed or food is processed, served or stored: In meat and food processing plants (processing areas) use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting station. Remove all baits and dead rodents before reuse of the plant (processing areas include storage and service).

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, DOMESTIC ANIMALS AND PETS. Place bait in locations protected at the time of application from access to children, domestic animals, pets and wildlife. Keep away from feed or foodstuffs. Harmful or fatal if swallowed or absorbed through the skin. This product should be handled with gloves. Do not get in the eyes, or the skin or on clothing. Wash arms, hands and face with soap and water after handling and before eating or smoking.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

FIRST AID

If in eyes, flush with plenty of water for 15 minutes. If on skin, wash thoroughly with soap and water. If swallowed, give patient one or two glasses of water and induce vomiting by touching back of the throat with a finger or blunt object. Repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Contact a physician or Poison Control Centre immediately.

TOXICOLOGICAL INFORMATION: If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. Vitamin K₁ in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case, measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

DISPOSAL

Dispose of unused, or unwanted product or dead rodents in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Do not reuse empty container. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted products and the cleanup of spills,

contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER

This control product is to be used only in accordance with the directions on the label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

(R)Registered trademark of Sanex Agro Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))