07-JAN-2009 2008-4587 Notification: Change in Registrant Address

## JAGUAR RODENTICIDE PELLET PLACE PACS

# KILLS RATS & MICE Kills Warfarin Resistant Norway Rats and House Mice

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

REGISTRATION NO.: 26129 PEST CONTROL PRODUCTS ACT

#### **DOMESTIC**

DANGER



**POISON** 

**GUARANTEE** 

Brodifacoum......0.0025%

Contains Denatonium Benzoate Warning, contains the allergen wheat. Contains malathion at 0.01%, as a preservative.

## READ THE LABEL AND OUTER PACKAGING BEFORE USING KEEP OUT OF REACH OF CHILDREN

For indoor use only

Notification Change

NET CONTENTS: 50g 3699 Kinsman Blvd., Madison WI 53704 USA

Manufactured by: Motomco Ltd., P.O. Box 8422, Madison, WI 53708 USA

Tel: 608-241-0202

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed though the skin. Do not open premeasured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately

from other laundry, with soap and water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS.

**FIRST AID:** If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and INDUCE VOMITING by touching back of throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or Poison Control Centre. If a pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin  $K_1$  in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**ENVIRONMENTAL HAZARD:** This product is toxic to fish, birds and wildlife. Keep out of lakes, rivers, streams or ponds.

**DIRECTIONS FOR USE:** Refer to exterior packaging for complete directions.

**DISPOSAL:** Do not reuse empty container. Dispose of empty container by placing in household garbage.

**Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**STORAGE:** Store in cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

031805

## JAGUAR RODENTICIDE PELLET PLACE PACS

## KILLS RATS & MICE Kills Warfarin Resistant Norway Rats and House Mice

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

REGISTRATION NO.: 26129 PEST CONTROL PRODUCTS ACT

#### **DOMESTIC**

**DANGER** 



**POISON** 

**GUARANTEE** 

Brodifacoum......0.0025% Contains Denatonium Benzoate

Warning, contains the allergen wheat. Contains malathion at 0.01%, as a preservative.

### READ THE LABEL BEFORE USING

(see back panel for additional precautionary statements) **KEEP OUT OF REACH OF CHILDREN** 

For indoor use only

Notification Change

NET CONTENTS: 40 x 50g 3699 Kinsman Blvd., Madison WI 53704 USA

Manufactured by: Motomco Ltd., P.O. Box 8422, Madison, WI 53708 USA

Tel: 608-241-0202

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed though the skin. Do not open premeasured place packs. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS.

**FIRST AID:** If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and INDUCE VOMITING by touching back of throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If a pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin  $K_1$  in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**ENVIRONMENTAL HAZARD:** This product is toxic to fish, birds and wildlife. Keep out of lakes, rivers, streams or ponds.

**DIRECTIONS FOR USE:** Use bait to control the Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*), and House mouse (*Mus musculus*) inside of buildings.

**USE LIMITATIONS:** Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Remove as much food as possible.

### **APPLICATION DIRECTIONS:**

Do not open or remove the plastic wrap from bait place pacs.

**RATS:** Place 2 to 8 Place Pacs at intervals of 4.5 to 9 metres per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

**HOUSE MICE:** Place 1 Place Pac at intervals of 2.5 to 3.5 metres per placement. Larger placements (up to 2 Place Pacs) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**RATS/MICE:** Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation

does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**DISPOSAL:** Do not reuse empty container. Dispose of empty container by placing in household garbage.

**Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**STORAGE:** Store in cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

\*\*\*\*\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.