

85.12.31

SANEX

GARDENER'S
WEED PREVENTER GRANULES

FOR VEGETABLE AND FLOWER GARDENS

DOMESTIC

NET CONTENTS 500 g

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

REG. NO. 18106
PEST CONTROL PRODUCTS ACT

Guarantee:
Chloramben 4.0%
(Present as Amonium Salts)

Manufactured by SANEX INC.
2695 Slough St., Mississauga, Ontario L4T 1G2
Branches throughout North America

DIRECTIONS: Sanex Gardener's Weed Preventer Granules kill weed seedlings as they sprout. It will maintain weed control for 6-8 weeks or more depending on weather conditions.

Apply to bare soil or to soil immediately after cultivation and weeding. It will not kill growing weeds. Do not disturb treated soil. Keep as much off the foliage as possible. Do not apply when foliage is wet.

If no rain falls within 3 to 4 days after application, water the area lightly to activate the chemical. If no water is available, a very shallow cultivation may be necessary to control emerged weeds.

For the second application later in the summer, it is often desirable to stir soil in flower and shrub beds to break up the crust that usually forms. Remove any weeds, stir the soil, then spread as recommended. A treatment at this time will prevent most weed growth for the rest of the season. Vegetable beds can be treated only once in a season.

This package will treat up to 75m². See individual plant or crop sections for application rates. Remove the container lid to expose a perforated cover and shake out contents avoiding existing plant foliage as much as possible.

Application Technique: Hold container in the middle and tilt top towards the ground at 45 degree angle. Rotate hand in a short diameter, circular motion at waist height. As the granules come out of the container move the container forward at the speed of a medium walk. If granules are spreading a little heavy, move forward at a faster pace. The width of the pattern given by this operation will be about 60 cm.

WEEDS CONTROLLED: Carpet weed, chickweed, lamb's quarters, pigweed, purslane, ragweed, smartweed, velvetleaf, barnyard grass, crab grass, foxtails, fall panicum. Sprouts from roots of established perennial weeds are not controlled.

ESTABLISHED PLANTS: Perennial flowers, shrubs and trees - Spread on the soil immediately after weeding and cleaning up each bed in the spring. Gladioli (flower production) - Plant corms 10 to 15 cm deep. Make two applications one at planting and one at lay-by after reshaping the bed (50 to 70 days after planting). Apply to give a cover of approximately 18 granules per 6.5 cm².

TRANSPLANTED PLANTS: Annual Flowers, Tomatoes and Peppers, Perennial Flowers and Shrubs - Prepare the bed and set out the transplants as usual. Allow about two weeks to overcome the "shock" of transplanting and start growing. Pull or hoe out any weeds that have started to grow. Spread granules evenly keeping as much off the foliage as possible. Apply to give a cover of approximately 18 granules per 6.5cm². Do not use on pansies.

SEEDED VEGETABLES: Seeding Asparagus, Carrots, Snap Beans (clay or organic soils only), Lima Beans, Squash and Pumpkin - Prepare a good seed bed, then plant the seeds. Spread granules the same day or within a few days of planting before weeds emerge. NOTE: Emerged seedling asparagus may be treated. The soil must be clean, cultivated and free of weeds before treating. Apply 9 to 14 granules per 6.5 cm² for asparagus, pumpkin and squash; 14-18 granules per 6.5 cm² for carrots; 14 granules per 6.5 cm² for lima beans; and 9 granules per 6.5 cm² for snap beans. Use the lower rates on light or sandy soils.

LAYBY TREATMENT FOR POTATOES: Apply granules after hilling when potatoes are 15 to 20 cm in height. Apply 14 to 18 granules per 6.5 cm². Use the lower rate on light or sandy soils.

LIST OF ANNUAL AND PERENNIAL FLOWERS, SHRUBS AND TREES which will not be injured when directions are followed. Arborvitae, ash, aster, azalea, celosia, chrysanthemum, dahlia, daylily, iris, English ivy, juniper, marigold, peony, petunia, privet, rhododendron, rose, snapdragon, spirea, yew, zinnia. Do not use on seeded flowers prior to emergence, as many will be injured.

LIMITATIONS:

1. Poor weed control and some crop damage may occur on sandy soils if heavy rains should wash the chemical into the soil.
2. Do not use on asparagus to be eaten the same year.

PRECAUTION: Avoid contact with skin, eyes and clothing. Avoid unintentional mixing with fertilizer, seeds, insecticides, fungicides or other herbicides. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate any body of water.

DISPOSE of empty container in garbage.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to person or property arising from the use or handling of this product and accepts the product on that condition.

This package is sold by weight not by volume. Packed as full as practicable by modern automatic equipment, it contains full net weight indicated. If it does not appear full when opened it is because contents have settled down during shipping and handling.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))