

96.04.02

SANEX(R)

TRI-KIL
SUPREME WEED KILLER

DOMESTIC

For Control of Hard-to-Kill Weed Species in Lawns

Controls: Bedstraw, Black Medick, (Yellow Clover), Buttercup, Chickweed, Clover, Dandelion, Devil's Paint Brush, English Daisy, Ground Ivy (Creeping Charlie), Heal-All, Knotweed, Narrow-leaved Plantain (Ribgrass), Poison Ivy, Ragweed, Shepherd's purse, Wild White Clover.

500 mL (1 L) (2 L) (4 L)

GUARANTEE:

2,4-D	95 g/L
Mecoprop (d-isomer)	50 g/L
Dicamba.....	9 g/L

(All present as amine salts)

READ THE LABEL BEFORE USING

REG. NO. 19395 P.C.P. ACT

SANEX AGRO INC.
36 Head Street
Dundas, Ontario
L9H 3H3

DIRECTIONS:

1 tsp. = 5 mL 1 tsp. = 15 mL

Mix 6 mL Tri-Kil per litre of water. Spray 20 litres of mixture evenly over 100 m². Apply in late spring or early fall on actively growing weeds. Avoid applications during long dry periods. Repeat as necessary. Fertilize and water lawn after weeds are killed, to fill any bare spots.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes and clothing. Protective clothing and equipment: Wear long-sleeved shirt, long pants, socks, shoes and chemical-resistant (e.g. rubber) gloves. Operator use precautions: Rinse gloves before removal. Wash hands before eating, drinking, using tobacco or using the toilet. If pesticide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing. Remove clothing and launder separately before reuse and promptly and thoroughly wash hands and exposed skin with soap and water, then shower. Reuse gloves for pesticide application only.

Avoid spray drift onto susceptible plants. Spray injurious to many cultivated species. Do not apply when rain is expected. Do not water within 24 hours of spraying. Do not use on Bent grasses. Do not use same sprayer for other uses. Do not apply on newly seeded lawns until after 3rd mowing. Do not store near seeds, feeds, fertilizers, insecticides or fungicides.

Reentry interval for turf: Do not allow people (other than the applicator) or pets on the treatment area during application. Do not enter treated areas until spray has thoroughly dried.

Maximum application frequency to turf: For good turf/lawn management, normally two applications per year per treatment site are adequate. This does not include spot treatments.

FIRST AID:

If swallowed, induce vomiting. Obtain prompt medical attention or contact a Poison Control Centre. If in eyes, flush with clear water for 15 minutes and get medical attention or contact a Poison Control Centre. If on skin, wash with soap and water.

DISPOSAL: Do not reuse container. Wrap in newspaper and dispose of empty container with household garbage.

(R) Registered trademark of Sanex Agro Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))