

85.12.31

Later's

Sevin(R) 40%

Liquid Insecticide

For Lawns, Ornamentals, Vegetables, and Fruits.

DANGER

POISON

DOMESTIC

READ THE LABEL BEFORE USING

REG. NO. 15775 P.C.P. ACT

Guarantee: Carbaryl 40%

250 mL

Later Chemicals Ltd.  
12080 Horseshoe Way,  
Richmond, B.C.  
V7A 4V5

SEVIN(R) is a registered trademark of Rhone Poulenc, Nederland, B.V.

1 Teaspoon = 5 mL  
1 Tablespoon = 15 mL

DIRECTIONS FOR USE:  
SHAKE WELL BEFORE USING  
DO NOT FREEZE.

Later's Sevin Liquid Insecticide is effective for the control of many insects on Lawns, Ornamental Plants, Vegetables and Fruits when used as recommended.

LAWNS: To control Chinch Bugs, Leaf Hopper, Fall Armyworm, Sod Webworm, Ants, Millipedes, Earwigs, Fleas and Mosquitoes. Mow lawn and apply at the rate of 25-30 mL with 15-20 L of water to cover 10 m<sup>2</sup> of area. For larger area, mix 250-300 mL with 150-200 L of water to cover 100 m<sup>2</sup> of area. Do not water for 2 days after treatment. Repeat in 2-3 weeks if necessary.

ORNAMENTALS: Arborvitae, Azalea, Birch, Boxwood, Carnation, Chrysanthemum,

Dogwood, Elm, Gladiolus, Holly, Hydrangea, Juniper, Lilac, Maple, Oak, Pines, Rose, Zinnia. To control Bagworm, Birch Leafminer, Black Vine Weevil, Blister Beetles, Boxelder Bug, Boxwood Leafminer, Cankerworms, Elm Leaf Aphid, Elm Leaf Beetles, Flea Beetles, Japanese Beetles, June Beetles, Lace Bugs, Leaf Hoppers, Leaf Rollers, Mealybugs, Pine Saw Flies, Plant Bugs, Psyllids, Rose Aphids, Rose Chafer, Rose Slugs, Scale Insects, Tent Caterpillars, Thrips (exposed), Willow Leaf Beetles: Use 300-400 mL per 100 L of water when insect or damage appears. Thoroughly spray the plant. Spray in early morning or late evening to avoid injury from high temperature. Repeat at intervals of 1-2 weeks, or as required. For Scale insects, apply when crawlers are present.

NOTES: Do not use on Boston Ivy or Virginia Creeper. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

NUMBER SHOWN AFTER EACH CROP INDICATES MINIMUM NUMBER OF DAYS BETWEEN LAST APPLICATION AND HARVEST.

Mix 20 mL per 10 L of water and spray to the point of run-off. Apply when insects or damage appears. Repeat at 7 - 10 day intervals or as necessary.		
FRUITS & BERRY	INSECTS	
Apples (7), Apricots(3), Peaches (1), Pears(7), Plum(1), Cherries(1) Strawberries(1)	Aphids, Apple maggots, Caterpillars, Codling moth, Fruit fly, Leafhoppers, Leafminers, Leaf rollers, Mealy bugs, Pear slug, Scale insects, Spider mites	
Mix amount shown with sufficient water for good coverage of 100 m2. Use lower rate on young plants. Apply when insects or damage appears. Repeat at 7 - 10 day intervals.		
VEGETABLES	INSECTS	AMOUNT
Beet root(7), Broccoli(3), Cabbage(3), Brussel sprout(3), Carrot(1), Celery(3), Cauliflower(3), Chinese cabbage(14), Leaf lettuce(14), Parsley(14), Radish(7), Spinach(14), Swiss chard(14), Turnips(14)	Flea Beetle, Leaf hopper	30 -75 mL
Cucumbers(3), Pumpkins(3), Squash(3)	Cucumber beetles, Climbing cutworms, Flea beetles, Leaf hoppers, Squash bugs	75 mL
Egg plant(1), Pepper(1), Tomato(1)	Colorado Potato beetles	30 mL
	Fall army worm, Tomato hornworm, Tomato fruit worm	75 mL

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed or inhaled. Do not breathe spray mist. Avoid skin contact. Wash hands before eating, drinking, or smoking. Shower or bathe after using. Wear regular long-sleeved work clothing. Avoid contamination of feeds, foods, feed troughs, and water receptacles. Do not contaminate streams, ponds and water supplies. Do not use where bees are foraging. Warn bee keepers to protect bees from treated area for one week after treatment. Highly toxic to bees. Some damage may occur to tender foliage in the presence of rain or high humidity of several days duration following spraying.

FIRST AID: Induce vomiting by sticking a finger down the throat or by giving warm water containing 15 gm (1 Tbsp.) of salt. If inhaled, move patient to hospital. Two 65 mg tablets of atropine may be given by mouth as an emergency measure.

TOXICOLOGICAL INFORMATION: Large doses of atropine may be needed to control symptoms.

DECONTAMINATION DISPOSAL: For small volumes of carbaryl suspended in water, add 1 kg of flake caustic per 2 1/2 kg of carbaryl, mix, and allow to stand 24 hours. Dispose of mixture through sewage treatment facility. Dispose of empty containers with household garbage.

Seller's guarantee shall be limited to the terms set out on this label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )