

19-DEC-1996

((LABEL))

KLOR (TM) 50E

LAWN, GARDEN AND HOME INSECTICIDE

EMULSIFIABLE CONCENTRATE

KEEP OUT OF REACH OF CHILDREN

DOMESTIC

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

DANGER POISON

GUARANTEE

Chlorpyrifos 50 g/L

REG. NO. 24625 P.C.P. ACT

Distributor:

United Horticultural Supply

R. R. #2

Dorchester, Ontario

NOL 1G5

NET: 250 mL

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid inhaling spray mist or contact with eyes, skin and clothing. May cause irritation to skin or eyes. Wash thoroughly after use. Avoid contamination of food, feed, utensils, drinking water or any body of water. Wash contaminated food surfaces, utensils and clothing with soap and water. DO NOT PERMIT CHILDREN OR PETS TO CONTACT SPRAYED AREAS UNTIL DRY. Do not use in poultry houses or other animal buildings. This product is toxic to fish and crustaceans.

FIRST AID: If swallowed, do NOT induce vomiting but rush patient to hospital, taking this labelled container with you. Contains petroleum distillate solvent. If on skin, wash with soap and water. Remove contaminated clothing. If in eyes, flush with plenty of water for 15 minutes and obtain medical attention.

TOXICOLOGICAL INFORMATION: Contains a cholinesterase inhibitor and a solvent. Atropine is an antidote.

DISPOSAL: Triple rinse container and add rinsings to mixture in sprayer. Wrap empty container in newspaper and dispose with household garbage. DO NOT RE-USE CONTAINER.

((LEAFLET))

KLOR (TM) 50E

LAWN, GARDEN AND HOME INSECTICIDE

EMULSIFIABLE CONCENTRATE

KEEP OUT OF REACH OF CHILDREN

DOMESTIC

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

DANGER POISON

GUARANTEE

Chlorpyrifos 50 g/L

REG. NO. 24625 P.C.P. ACT

Distributor:  
United Horticultural Supply  
R. R. #2  
Dorchester, Ontario  
NOL 1G5

NET: 250 mL

KLOR is an effective insecticide for vegetables, lawns, soil, ornamentals, houseplants and household use.

DIRECTIONS

5 mL = 1 teaspoon                      15 mL = 1 tablespoon

VEGETABLES: When damage from cutworms is evident, apply spray to give uniform coverage of soil area. Use 60 mL with 10 L of water per 100 m<sup>2</sup> on broccoli, Brussels sprouts, cabbage, Chinese cabbage, cauliflower, celery, cucumber, green pepper, sweet corn tomato. Make no more than 1 application per season. (See table below for days to wait between application and harvest). For carrots and onions, use 120 mL with 10 L of water and apply 3 L per 100 m of row of soil surface. Do not use on green bunching onions. For strawberries, use 120 mL in 10 L of water per 100 m<sup>2</sup> and apply to soil once between June 1 and June 15. For control of potato flea beetle and tarnished plant bug on potato apply 80 mL

in 10 L of water per 100 m<sup>2</sup>. Apply weekly as a foliar application as needed. Do not make more than 9 applications in one season on potatoes.

DAYS TO WAIT BETWEEN APPLICATION AND HARVEST:

Potato	7 days
Strawberries	20 days
Broccoli, Brussels sprouts, Cabbage,	
Chinese cabbage, Cauliflower	32 days
Green Pepper	40 days
Cucumber, Carrot, Onion	60 days
Sweet Corn, Celery, Tomato	70 days

LAWNS: USE NO MORE THAN 2 APPLICATIONS PER SEASON.

Apply when insects first appear and repeat when needed. The lawn surface should be moist before treatment. To control White Grubs and the grubs of the European Chafer and Japanese Beetle, mix 400 mL with 10 L of water per 100 m<sup>2</sup> of lawn area. Spray in late July or August when grubs are small and are feeding on the grass roots just below the soil surface. Immediately after spraying, water the lawn thoroughly. To control chinch bugs, sod webworms, ants and mosquitoes mix 200 mL with 10 L of water per 100 m<sup>2</sup> of lawn area (a watering can is suitable for this treatment). For Ants, spray nests and surrounding areas. For Sod Webworms, delay watering or mowing for 12 - 24 hours after treatment. To control Crane Fly Larvae, (Leatherjackets) use 200 mL with 10 L of water per 100 m<sup>2</sup>. Apply as a drenching spray in water in the late fall after the flight of Adult crane flies has ceased for the year.

ORNAMENTAL PLANTS AND SHRUBS: Make no more than 2 treatments per season. Soil treatment - To control Ants, Crickets, Earwigs, Sowbugs, Spiders, Cutworms, White Grubs, European Chafer Larvae, and Japanese Beetle Larvae, spray soil and turf around plants to give good coverage. Apply 200 mL with 10 L of water per 100 m<sup>2</sup>.

ORNAMENTAL PLANTS, SHRUBS, AND TREES: Foliar treatment - To control Aphids, Ash and Lilac Borers, Mealy Bugs, Mites, Scales, Spittle Bugs, Thrips and Whiteflies, on deciduous and coniferous trees and shrubs, mix 200 mL with 10 L of water and thoroughly spray all foliage. NOTE: DO NOT USE ON ROSES, AZALEAS, CAMELLIAS, POINSETTIAS, GERANIUMS, OXALIS, COLEUS, OR VARIEGATED IVY.

NATIVE ELM BARK BEETLE: To prevent the adult beetle from overwintering in uninfected trees and to reduce beetle populations in disease free areas, mix 300 mL with 3 L of water and spray the bottom 2.5 m of trunk. Wet the trunk thoroughly but do not spray to runoff. Make only one application per season. To prevent branch and twig feeding on uninfected trees, spray the tree crown thoroughly. Also spray the base of the tree where the roots spread. For best results, consult municipal authorities to incorporate a complete integrated Dutch Elm disease program.

INSECT PESTS INDOORS: For Cockroaches, Ants, Crickets, Earwigs,

Firebrats, Fleas, Silverfish, Sowbugs, Spiders, Brown Dog Ticks, mix 50 mL with 1 L of water and apply as a spot treatment using a coarse spray or paint brush on surfaces, along baseboards, and behind fixtures where these pests are a problem. For Fleas and Ticks, mix 100 mL with 1 L of water and apply spray to pet beds and resting quarters, as well as nearby cracks and crevices. Replace bedding after treatment. DO NOT TREAT PETS. For Carpet Beetles, mix 100 mL with 1 L of water and apply along baseboards, under carpets and wherever insects are seen. Do not spray bedding, clothing, or furniture. Repeat if reinfestation occurs, but no more than once every 2 months. DO NOT USE AS A SPACE SPRAY. Cover or remove all food before treatment. Do not allow spray to contact food, food preparing surfaces, utensils or drinking water.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid inhaling spray mist or contact with eyes, skin and clothing. May cause irritation to skin or eyes. Wash thoroughly after use. Avoid contamination of food, feed, utensils, drinking water or any body of water. Wash contaminated food surfaces, utensils and clothing with soap and water. DO NOT PERMIT CHILDREN OR PETS TO CONTACT SPRAYED AREAS UNTIL DRY. Do not use in poultry houses or other animal buildings. This product is toxic to fish and crustaceans.

FIRST AID: If swallowed, do NOT induce vomiting but rush patient to hospital, taking this labelled container with you. Contains petroleum distillate solvent. If on skin, wash with soap and water. Remove contaminated clothing. If in eyes, flush with plenty of water for 15 minutes and obtain medical attention.

TOXICOLOGICAL INFORMATION: Contains a cholinesterase inhibitor and a solvent. Atropine is an antidote.

DISPOSAL: Triple rinse container and add rinsings to mixture in sprayer. Wrap empty container in newspaper and dispose with household garbage. DO NOT RE-USE CONTAINER.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

