

90.04.18

STARBAR

INSECTICIDE

Louse Powder

AGRICULTURAL

GUARANTEE: Carbaryl..... 5 %

REGISTRATION NO: 14268 PEST CONTROL PRODUCTS ACT

CONTROLS

Biting and sucking lice, fleas, winter ticks and horn flies on:
Dairy and beef cattle, horses, hogs and sheep, Northern Fowl
mites, chicken mites, fleas, lice, fowl ticks and bed bugs in
Poultry and Poultry houses.

KEEP OUT OF REACH OF CHILDREN
READ THE LABEL BEFORE USING
STORE PRODUCT IN ORIGINAL CONTAINER
KEEP TIGHTLY CLOSED AND IN A COOL, DRY PLACE

750 g

Zoecon Canada Inc.
12 Stanley Court, Unit 3
Whitby, ONTARIO L1N 8P9
PHONE: (416) 430-2511

Starbar is a Registered Trademark of Sandoz Ltd.

DIRECTIONS

In or around poultry houses and farm buildings: For the control
of Northern Fowl and Chicken Mites, Fleas and Bed Bugs, apply
dust thoroughly to floor, litter and adjacent areas at the rate
of 1.25 Kg per 10 sq. metres of surface. Repeat as necessary.

On the birds: For the control of Northern Fowl and Chicken Mites, Lice, Fleas and Fowl Tick, apply 500 g of dust per 100 birds. Repeat in 4 weeks or as needed. Apply dust using a hand, rotary or puff duster.

Caution: Do not use around or on birds within 7 days of slaughter. Do not treat nests and avoid direct contamination of eggs. Avoid contamination of feed and water troughs.

On beef cattle, horses, sheep and hogs: For the control of Winter Ticks, Lice, Fleas and Horn Flies, dust entire animal thoroughly using not more than 500 g of dust per head. Repeat if necessary, but not more than twice weekly. For louse control repeat in 14 to 18 days.

On dairy cattle: Use as for Beef Cattle but DO NOT APPLY DIRECTLY TO THE UNDERLINE PORTION AND THE UDDER. Repeat if necessary, but not more often than twice weekly. DO NOT APPLY BEFORE MILKING: ONLY IMMEDIATELY AFTER MILKING WASH UDDER THOROUGHLY BEFORE MILKING. DO NOT CONTAMINATE MILKING UTENSILS.

Caution: Do not apply to livestock within one week of slaughter. Do not exceed dosage given above.

Precautions: KEEP OUT OF REACH OF CHILDREN. Harmful if inhaled or swallowed. Do not take internally. Avoid unnecessary skin contact. Wash hands before eating. Take a shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Avoid contamination of feed, foodstuffs, feeding trough and watering receptacles.

Toxic to bees: Do not use where bees are foraging.

First Aid: If swallowed, induce vomiting and get medical aid. If in eyes, flush thoroughly with plenty of water.

TOXICOLOGICAL INFORMATION: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal. Contact physician at Poison Control Centre.

DISPOSAL: Empty container thoroughly. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product and the clean up of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))