30-JAN-2003. Notification of Label Changes received. See below for label changes.

07-SEP-1999

((LABEL))

#### 2 CHAMPION®

#### HERBICIDE

#### EMULSIFIABLE CONCENTRATE

THIS PRODUCT MUST BE USED IN COMBINATION WITH EXTRA™ HERBICIDE OF THE CHAMPION EXTRA HERBICIDE TANK MIX

FOR USE IN BARLEY

PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA

#### COMMERCIAL

CAUTION POISON

#### TRIPLE RINSE

DANGER - EYE AND SKIN IRRITANT

Equipment may be required

REGISTRATION NO. 25538 PEST CONTROL PRODUCTS ACT

GUARANTEE: fenoxaprop-p-ethyl 92 g/L

DISTRIBUTED BY: DuPont Canada Inc. Agricultural Products Box 2300, Streetsville Mississauga, ON L5M 2J4 Telephone: 1-800-667-3925

# Notification Change

Champion Herbicide is manufactured and registered by: Bayer AgrEvo Canada Inc. Suite 1

295 Henderson Drive

Bayer Cropscience Inc. Suite 100 3131-114th Ave. S.E. Calgary, AB T2Z 3X2

# Regina, SK S4N 6C2 Telephone: 1-800-667-5959

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

NET CONTENTS: CHAMPION 8.1 LITRES

#### PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. DANGER - Eye and skin irritant causes eye damage. Protective eye equipment must be worn. Do not get in eyes, on skin or clothing. Harmful if inhaled. Do not breathe vapour or spray mist. Protective Clothing and Equipment: Wear coveralls over normal work clothes during all activities. Wear chemical resistant gloves and apron, eye protection (goggles or faceshield) and respirator during mixing, loading, repairs and cleanup. Change contaminated clothing daily and wash before reusing. DO NOT handle product with bare hands. Chemical resistant gloves significantly reduce hand exposure. ALWAYS wear gloves for mixing/loading operations and when making sprayer and nozzle repairs and adjustments. DO NOT USE LEATHER OR CLOTH GLOVES. Rinse gloves with soap and water before removal and wash hands thoroughly before eating, smoking or drinking. Clean protective equipment daily. FIRST AID - INGESTION: If the tank-mix spray solution or CHAMPION Herbicide is swallowed do NOT induce vomiting. Immediately contact a poison control centre or physician. EYE CONTACT: If in contact with eyes, flush with plenty of water for 15 minutes. Immediately contact a poison control centre or physician. SKIN CONTACT: In case of skin contact, remove contaminated clothing and wash skin thoroughly with soap and water. If irritation persists, contact a physician. INHALATION: If inhaled, transfer individual from the site of exposure to fresh air. TOXICOLOGICAL INFORMATION: Medical Personnel should contact AgrEvo's Medical Information Services: 1-800-228-5635, ext. 211. CHAMPION contains PETROLEUM DISTILLATES. ENVIRONMENTAL PRECAUTIONS AND INFORMATION: CAUTION: This product is toxic to fish, daphnids, aquatic plants, terrestrial plants and some terrestrial invertebrates. Overspray or drift to aquatic bodies of water, such as ponds, lakes, rivers, streams, irrigation water and terrestrial wildlife habitats such as wetlands, wood lots, shelterbelts, sloughs or dry slough borders, vegetated areas at the edges of bodies of water and other terrestrial wildlife habitats should be avoided. Leave a buffer zone of 15 metres between the last spray swath and the edge of any of these habitats. Avoid spray drift to susceptible plants and non-target areas. Do not contaminate

irrigation water or apply to land where tail water will be used to irrigate crop land. USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES. STORAGE: Keep away from fire or open flame or other sources of heat. CANNOT be stored below freezing. If stored for 1 year or longer, shake well before using. Store the tightly closed container away from feeds, seeds, fertiliser, plants and foodstuffs. Do not use or store in or around the home. Keep in original container during storage. CONTAINER DISPOSAL: Triple rinse the emptied containers thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the containers prior to their disposal. Make the empty containers unsuitable for further use. Dispose of the containers in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer. NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

CHAMPION is manufactured and registered by AgrEvo Canada Inc. CHAMPION EXTRA is distributed by DuPont CANADA INC. CHAMPION EXTRA is a trademark of DuPont Canada Inc. [duPont logo] and CHAMPION are registered trademarks of E.I. duPont de Nemours and Company.

07-SEP-1999

((DIRECTIONS FOR USE))

DUPONT

CHAMPION EXTRA

#### HERBICIDE TANK MIX

#### POST-EMERGENT

#### FOR USE IN BARLEY

# PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA

#### COMMERCIAL

# CAUTION POISON

#### TRIPLE RINSE

#### DANGER - EYE AND SKIN IRRITANT

Equipment may be required

THIS TANK MIX USES: CHAMPION Herbicide Registration No. 25538 PEST CONTROL PRODUCTS ACT GUARANTEE: fenoxaprop-p-ethyl 92 g/L

EXTRA Herbicide Registration No. 25537

PEST CONTROL PRODUCTS ACT GUARANTEE: thifensulfuron methyl 50% tribenuron methyl 25%

DISTRIBUTED BY: DuPont Canada Inc. Agricultural Products Box 2300, Streetsville Mississauga, ON L5M 2J4 Telephone: 1-800-667-3925

Champion Herbicide is manufactured

Notification Change

and registered by: AgrEvo Canada Inc. 295 Henderson Drive Regina, SK S4N 6C2 Telephone: 1-800-667-5959

Bayer Cropscience Inc. Suite 100 3131-114th Ave. S.E. Calgary, AB T2Z 3X2

READ THE LABEL BEFORE USING

NET CONTENTS: CHAMPION - 8.1 LITRES EXTRA - 162 GRAMS

CHAMPION EXTRA HERBICIDE TANK MIX

The Product

Table of Contents:

Section Number:

GENERAL INFORMATION

SAFETY AND HANDLING

Precautions, Protective

1

	Clothing and Equipment First Aid and Toxicological	2
	Information	3
	Environmental Precautions and Information	4
	Storage	5
	Disposal	б
DIRECTIONS FOR USE	Crops, Weeds, Rates, Timing and	
	Cautions	7
	Tank Mixes	8
	Application Instructions, Cautions and	
	Rainfast Period	9
	Mixing Instructions	10
	Sprayer Cleanup	11
	Herbicide Resistance Management	12
	Livestock Feeding and Pre-Harvest	
	Intervals	13
	Notices	14

# For More Information, Contact:

DUPONT CANADA INC. AGRICULTURAL PRODUCTS Box 2300, Streetsville Mississauga, ON L5M 2J4 **Telephone:** 1-800-667-3925

GENERAL INFORMATION

Section 1: The Product

FOR USE IN THE PRAIRIE PROVINCES AND PEACE RIVER REGION OF BRITISH COLUMBIA.

CHAMPION EXTRA Herbicide Tank Mix provides control of certain annual grassy and broadleaf weeds in Barley.

One 8.1 litre container of the CHAMPION Herbicide component, mixed with one 162 gram container of the EXTRA Herbicide component will treat 8.1 hectares (20 acres).

CHAMPION EXTRA Herbicide Tank Mix may NOT be used on any crop for which it is NOT REGISTERED.

SAFETY AND HANDLING

#### Section 2: Precautions, Protective Clothing and Equipment

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

DANGER - EYE AND SKIN IRRITANT CAUSES EYE DAMAGE. Protective eye equipment must be worn.

Do not get in eyes, on skin or clothing.

Harmful if inhaled. Do not breathe vapour or spray mist.

Protective Clothing and Equipment:

Wear coveralls over normal work clothes during all activities. Wear chemical resistant gloves and apron, eye protection (goggles or faceshield) and respirator during mixing, loading, repairs and cleanup.

Change contaminated clothing daily and wash before reusing.

DO NOT handle product with bare hands. Chemical resistant gloves significantly reduce hand exposure. ALWAYS wear gloves for mixing/loading operations and when making sprayer and nozzle repairs and adjustments. DO NOT USE LEATHER OR CLOTH GLOVES.

Rinse gloves with soap and water before removal and wash hands thoroughly before eating, smoking or drinking.

Clean protective equipment daily.

#### Section 3: First Aid and Toxicological Information

#### FIRST AID:

INGESTION: If the tank-mix spray solution or CHAMPION Herbicide is swallowed do NOT induce vomiting. Immediately contact a poison control centre or physician.

If EXTRA Herbicide is swallowed, administer water freely and induce vomiting. Repeat until vomit fluid is clear. Contact a poison control centre or physician.

EYE CONTACT: If in contact with eyes, flush with plenty of water for 15 minutes. Immediately contact a poison control centre or physician.

SKIN CONTACT: In case of skin contact, remove contaminated clothing

and wash skin thoroughly with soap and water. If irritation persists, contact a physician.

INHALATION: If inhaled, transfer individual from the site of exposure to fresh air.

TOXICOLOGICAL INFORMATION:

Medical Personnel should contact AgrEvo's Medical Information Services toll-free: 1-800-228-5635, ext. 211.

The CHAMPION Herbicide component of this tank mix contains PETROLEUM DISTILLATES.

#### Section 4: Environmental Precautions and Information

CAUTION: This product is toxic to fish, daphnids, aquatic plants, terrestrial plants and some terrestrial invertebrates. Overspray or drift to aquatic bodies of water, such as ponds, lakes, rivers, streams, irrigation water and terrestrial wildlife habitats such as wetlands, wood lots, shelterbelts, sloughs or dry slough borders, vegetated areas at the edges of bodies of water and other terrestrial wildlife habitats should be avoided. Leave a buffer zone of 15 metres between the last spray swath and the edge of any of these habitats.

Avoid spray drift to susceptible plants and non-target areas.

Do not contaminate irrigation water or apply to land where tail water will be used to irrigate crop land.

USE ONLY FOR RECOMMENDED PURPOSES AND AT RECOMMENDED RATES.

#### Section 5: Storage

Keep away from fire or open flame or other sources of heat.

CANNOT be stored below freezing.

If stored for 1 year or longer, shake well before using.

Store the tightly closed container away from feeds, seeds, fertilizers, plants and foodstuffs.

Do not use or store in or around the home.

Keep in original container during storage.

#### Section 6: Disposal

WASTE DISPOSAL

If any spray solution remains in the tank after spraying is finished, it should be sprayed on the perimeter of the field just sprayed, NOT ON THE CROP, away from water supplies, ditches, irrigation canals and desirable vegetation.

#### CONTAINER DISPOSAL

1. Triple rinse the emptied containers thoroughly and add the rinsings to the spray mixture in the tank.

- 2. Follow provincial instructions for any required additional cleaning of the containers prior to their disposal.
- 3. Make the empty containers unsuitable for further use.
- 4. Dispose of the containers in accordance with provincial requirements.
- 5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer.

# DIRECTIONS FOR USE

# Section 7: Crops, Weeds, Rates, Timing and Cautions

CROP	WEED	RECOMMENDED TIME OF APPLICATION	APPLICATION RATE
BARLEY	ANNUAL GRASSY WEEDS: Wild Oats Green Foxtail Barnyard Grass	1-6 leaves (up to emergence of the 3rd tiller)	Apply CHAMPION at 1 L/ha. Apply EXTRA at 20 g/ha
	BROADLEAF WEEDS: Annual Smartweed (Green Smartweed, Lady's-thumb) Wild Mustard Corn Spurry Cow Cockle Hemp-nettle Kochia Lamb's-quarters Redroot Pigweed Russian Thistle Shepherd's-purse Stinkweed Tartary Buckwheat Volunteer Canola 1, Wild Buckwheat 2, Chickweed 3	less than 10 cm tall or across Cotyledon to 3 weed leaf stage 1-6 weed leaf stage	Apply CHAMPION at 1 L/ha. Apply EXTRA at 20 g/ha
	SUPPRESSION ONLY: Canada Thistle 4	up to 15 cm height	

Cleavers	1-3 whorls	
Round-leaved Mallow 5	2-6 weed leaf stage	
Scentless Chamomile	up to 10 cm tall or across	
Sow Thistle 4	up to 15 cm height	
Stork's-bill	2-6 weed leaf stage	
Toadflax 6	up to 15 cm height	

# NOTES:

- 1 **Volunteer Canola:** CHAMPION EXTRA Herbicide Tank Mix alone will not control imazethapyr tolerant Canola varieties, e.g. Canola varieties with the Pursuit Smart® trait.
- 2 Wild Buckwheat: Apply CHAMPION EXTRA Herbicide Tank Mix to actively growing Wild Buckwheat in the cotyledon to 1-3 leaf stage. Under stress conditions, such as dry weather, control may be reduced. Large plants may regrow after treatment.
- 3 **Chickweed:** Apply CHAMPION EXTRA Herbicide Tank Mix when the Chickweed is small (1-6 leaf stage) and actively growing but before crop canopy prevents thorough coverage of weeds. Chickweed emerging after application will not be controlled.
- 4 **Canada Thistle, Sow Thistle:** Apply CHAMPION EXTRA Herbicide Tank Mix components when the majority of the thistles have emerged and are actively growing. For best top growth control, apply when thistles are no larger than 15 cm tall, before bud stage, and before crop canopy prevents thorough coverage of thistles. A single application will effectively inhibit the ability of emerged thistles to compete with the crop. Later emerging thistle will not be controlled.
- 5 Round-leaved Mallow: Apply CHAMPION EXTRA Herbicide Tank Mix post-emergence when the actively growing Round-leaved Mallow is in the 2-6 leaf stage (10-12 cm in height). CHAMPION EXTRA Herbicide Tank Mix will keep mallow stunted, but may not reduce the overall populations.

6 **Toadflax:** Apply CHAMPION EXTRA Herbicide Tank Mix when Toadflax is not taller than 15 cm in height. A control program for this weed includes both frequent tillage and chemical application.

NOTE: Weed suppression is a visual reduction in weed competition as compared to the untreated check. Suppression can be expressed as reduced population, vigour and/or height and/or a yellowing or browning of the leaves. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

# CAUTION:

This Herbicide Tank Mix is registered for use in Barley only. Do not treat Barley if underseeded to Forage Crops.

Initial crop injury, such as discolouration, may be observed after application. This condition is temporary and should not affect yield. Early application is important to maintain crop safety.

#### Timing:

Crop damage will occur if application is made earlier or later than the leaf stages recommended on this label.

Treatment at the 3-4 leaf stage of crop and grassy weeds will combine maximum crop tolerance and weed susceptibility.

However, CHAMPION EXTRA Herbicide Tank Mix may be applied to Barley at the 2-5 leaf stage on the main stem with no more than 2 tillers.

Application of CHAMPION EXTRA Herbicide Tank Mix past the emergence of the second tiller will result in crop damage. The 2-tiller stage of Barley usually occurs within 25 days after seeding.

Apply to emerged, actively growing weeds according to the stages listed.

Weeds that emerge after application will not be controlled.

Some broadleaf weeds may not be controlled if infestation is heavy, weeds are in bud, or weather is dry and cool.

Under stressed conditions and/or heavy crop canopy, earlier application will result in improved grassy weed control.

# ((BARLEY STAGING ILLUSTRATION))

SECTION 8: TANK MIXES

For control of Wild Oats, Green Foxtail, Barnyard Grass and broadleaf weeds, CHAMPION EXTRA Herbicide Tank Mix can be tank mixed with MCPA Ester as instructed in the following table. Always read and follow the use directions, environmental restrictions and use precautions on the CHAMPION Herbicide and EXTRA Herbicide labels, and the label of the product being applied with CHAMPION EXTRA Herbicide Tank Mix.

Do not tank mix any other chemical additives, pesticides or fertilizers unless recommended on this label.

Product Name	Rate of Application	
CHAMPION EXTRA CHAMPION + EXTRA + MCPA Ester 500	1 L/ha + 20 g/ha + 840 mL/ha	
USE ONLY MCPA ESTER 500 in this tank mix and only at recommended rates. Failure to do so may result in reduced Wild Oat control. DO NOT USE MCPA AMINE.		

#### Section 9: Application Instructions, Cautions and Rainfast Period

#### Application Instructions:

GROUND APPLICATION ONLY. DO NOT APPLY BY AIR.

All the components in the CHAMPION EXTRA Herbicide Tank Mix package must be applied together according to label recommendations or a reduction in weed control and/or crop tolerance will occur.

Apply CHAMPION EXTRA Herbicide Tank Mix in a minimum of 110 L/ha of water at a pressure of 275 kPa, or 310 kPa if using check valves, and at ground speed of 8-10 km/h.

The use of 80E or 110E flat fan nozzles is recommended for optimum spray coverage.

Do not use flood jet nozzles, controlled droplet application equipment or Sprafoil® equipment.

Application of the spray at a 45E angle forward and the higher water volumes will result in better spray coverage and penetration of the crop canopy.

Uniform, thorough spray coverage is important to achieve consistent weed control.

Avoid overlapping; shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over-application .

Application of CHAMPION EXTRA Herbicide Tank Mix must take place within the leaf stages outlined for all weeds on this label. Late application will result in reduced weed control.

During periods of stress, plants are not actively growing. When daytime temperatures before or following application are very hot combined with very dry conditions and low humidity, plants are under stress. Application of CHAMPION EXTRA Herbicide Tank Mix during these periods may result in substantially reduced control.

Do not apply to crop that is stressed by severe weather conditions, frost, low fertility, drought, water saturated soil, disease or insect damage as crop injury may result.

#### Application Cautions:

DO NOT exceed a total of 20 g/ha per year of EXTRA Herbicide (thifensulfuron-methyl and tribenuron methyl).

DO NOT apply CHAMPION EXTRA Herbicide Tank Mix more than once a year.

EXTRA Herbicide must be tank mixed at the recommended rate with CHAMPION Herbicide or crop damage and/or a reduction in grassy and broadleaf weed control will result.

A time interval of 7 days prior to application, or 4 days after application of CHAMPION EXTRA Herbicide Tank Mix is required before any other pesticide can be applied.

Do not apply to terrain where there is a potential for surface runoff to enter aquatic systems.

Do not apply when environmental conditions such as wind speed, temperature, relative humidity, etc., are favourable for drift to occur.

# Rainfast Period:

Rainfall within 4 hours of application will result in reduced weed control of CHAMPION EXTRA Herbicide Tank Mix.

#### Section 10: Mixing Instructions

- 1. Ensure that the spray tank, lines and filter are thoroughly clean.
- 2. Fill the spray tank half full with clean water and start agitation or bypass system.
- 3. Add the required amount of EXTRA Herbicide (container #1) while maintaining agitation until EXTRA Herbicide is completely in suspension. Add all containers of EXTRA Herbicide if using more than one.
- Slowly add the correct amount of CHAMPION Herbicide (container #2) and continue agitation. Use all containers of CHAMPION Herbicide, if using more than one container.
- 5. Triple rinse the emptied containers into the spray tank.
- 6. Add the remaining amount of water while agitation continues.
- 7. On repeat loads, ensure that the amount of spray solution left in the spray tank from the previous load is less than 10% of the volume about to be mixed.
- 8. Spray out immediately. Spray mixture should not be left in the tank overnight. If spray solution is left standing without agitation, thoroughly agitate and re-suspend any settled CHAMPION EXTRA Herbicide Tank Mix before resuming the spraying operation.
- DO NOT tank mix with any other chemical additives, pesticides, surfactants or fertilizers.

## Section 11: Sprayer Cleanup

Thoroughly clean the spray tank by flushing with water/detergent mixture after using CHAMPION EXTRA Herbicide Tank Mix and before using any other pesticide.

The cleaning agent may be either household ammonia (minimum 3% ammonia) or an ammonia based cleaning agent containing at least 3% ammonia.

DO NOT use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which may cause eye, throat and lung irritation.

Do not drain or clean equipment or containers near well, water source or sensitive vegetation.

#### Cleanup Before You Spray:

IN ALL CASES, prior to spraying crops sensitive to CHAMPION EXTRA Herbicide Tank Mix, complete a thorough cleaning of the spray tank, lines and filter.

Spray equipment should be triple rinsed using a strong detergent solution. A scrubbing of the tank is recommended.

Follow the cleanup instructions below to ensure adequate sprayer cleaning and removal of the CHAMPION EXTRA Herbicide Tank Mix.

#### Cleanup After You Spray:

To avoid subsequent injury to other crops, immediately after spraying Barley and before spraying other crops, thoroughly remove all traces of CHAMPION EXTRA Herbicide Tank Mix from the mixing and spray equipment by flushing with a water/detergent mixture.

Drain or flush equipment and containers away from wells, water sources, and sensitive vegetation.

SPRAYER CLEANOUT INSTRUCTIONS WHEN MOVING INTO WHEAT, RYE OR FLAX

When moving into WHEAT, SPRING or FALL RYE, or FLAX immediately following the application of CHAMPION EXTRA Herbicide tank mix, clean the sprayer by thoroughly flushing with water/detergent or cleaning agent.

SPRAYER CLEANOUT INSTRUCTIONS WHEN MOVING INTO BROADLEAF CROPS:

Broadleaf crops can be damaged by CHAMPION EXTRA Herbicide Tank Mix residues in the spray tank even after a number of applications of a different product. It is critical to thoroughly clean and remove all traces of CHAMPION EXTRA Herbicide tank mix from the spray tank before moving into a broadleaf crop.

IN ALL CASES, before spraying a broadleaf crop (such as Canola, Peas, Lentils, Alfalfa, Sugar Beets, Vegetables, etc.) thoroughly clean the spray tank, as the EXTRA Herbicide component of the CHAMPION EXTRA tank mix can cause crop injury to sensitive crops at very low concentrations. Observe the following cleanup instructions to ensure satisfactory sprayer cleaning and removal of the CHAMPION EXTRA Herbicide Tank Mix.

CLEANUP INSTRUCTIONS PRIOR TO SPRAYING BROADLEAF CROPS:

1. Remove nozzles and screens and clean separately in a bucket containing water and cleaning agent.

2. Drain tank and flush tank, boom, and hoses with clean water for at least 10 minutes. Inspect tank to ensure all visible residues have been removed. If necessary, repeat this step.

# If the sprayer tank is equipped with rotary washing nozzles, go to step 6.

- 3. Fill the tank with clean water while adding 1 litre cleaning agent per 100 litres water. Flush solution through boom and hoses and then add sufficient water to fill the tank. Allow to stand for 15 minutes with agitation. Flush boom and hoses again, and drain tank.
- 4. Repeat step 3.
- 5. Go to and complete step 9.
- 6. If the sprayer tank is equipped with rotary nozzles, fill approximately one quarter full with water, and add 1 litre of cleaning agent for each 100 litres of water.
- 7. Operate the rotary nozzles for at least 15 minutes, ensuring that all inside surfaces of the tank are covered by the water/cleaning agent mixture. After about 5 minutes of agitation, flush the solution through the hoses and boom for about 2 minutes. Drain tank.
- 8. Repeat steps 6 and 7.
- 9. Thoroughly rinse the tank, hoses and boom with clean water for at least 5 minutes.

#### Section 12: Herbicide Resistance Management

Some naturally occurring biotypes of certain grassy weeds listed on this label have been identified in western Canada that are resistant to the CHAMPION Herbicide component of this herbicide tank mix and related products (those with the same mode of action). Some naturally occurring biotypes of certain broadleaf weeds listed on this label have been identified in western Canada that are resistant to the EXTRA Herbicide component of this herbicide tank mix and related products (those with the same mode of action). These resistant populations are able to survive a use rate several times higher than needed to control susceptible populations.

To help prevent a build-up of herbicide resistant weed populations, CHAMPION EXTRA Herbicide Tank Mix should be rotated with herbicides that have a different mode of action, or choose alternate cultural practices that are effective in controlling these weeds, i.e. Summerfallow, Chemfallow, preseeding burnoff, crop rotation, etc.

If poor performance cannot be attributed to adverse weather conditions or improper application methods, a resistant biotype may be present.

If resistance is suspected, contact your local representative of DuPont Canada Inc. for assistance. For more information, contact: DuPont Canada Inc. at 1-800-667-3925.

If resistance is identified, additional treatments with CHAMPION EXTRA Herbicide Tank Mix or related herbicides will be ineffective and is not recommended.

#### Section 13: Livestock Feeding and Pre-Harvest Intervals

Do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

Do not apply CHAMPION EXTRA Herbicide Tank Mix within 65 days of harvest.

#### Section 14: Notices

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FOR MORE INFORMATION CONTACT:

DU PONT CANADA INC. AGRICULTURAL PRODUCTS Box 2300, Streetsville Mississauga, ON L5M 2J4 Telephone: 1-800-667-3925

CHAMPION is manufactured and registered by AgrEvo Canada Inc.

CHAMPION EXTRA is distributed by DuPont CANADA INC.

CHAMPION EXTRA™ is a trademark of DuPont Canada Inc.

DUPONT LOGO and CHAMPION® are registered trademarks of E.I. duPont de Nemours and Company.

Pursuit and Smart are registered trademarks of Cyanamid Crop Protection Ltd.

SPRAFOIL is a Registered Trademark of D&W Industries.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))