

97/2002

Net Contents 4.5 L

CCS 06-3570-01-000

Proquat\*

**GERMICIDAL DETERGENT**

Germicidal, Fungicidal, Virucidal based on a unique blend of Quaternaries.

Recommended for the cleaning and disinfecting of vehicles, livestock and poultry buildings and equipment, hatcheries and equipment.

**GUARANTEE:** Didecyl Dimethyl Ammonium Chloride 1.4%, Alkyl (C12, 40%; C14, 50%; C16, 10%) Benzyl Dimethyl Ammonium Chloride 11.2%.

**AGRICULTURAL**

**DANGER**

**CORROSIVE & POISON**

REGISTRATION NO: 15501

PEST CONTROL PRODUCTS ACT

**READ THE LABEL BEFORE USING**

Upjohn  
**Pharmacia & Upjohn Animal Health**  
40 Centennial Road  
Orangeville, Ontario, L9W 3T3

PROQUAT Germicidal Detergent is a concentrated germicide based on a unique blend of quaternaries. PROQUAT cleans, disinfects and deodorizes in soft or hard water (by official A.O.A.C. test 400 ppm CaCO<sub>3</sub>). For heavily soiled areas, a preliminary cleaning with PROQUAT is necessary before final disinfection.

**Germicidal**      Staphylocidal Pseudomonacidal  
A.O.A.C. Use dilution confirmation.....1:256

**Fungicidal** T. mentagrophytes A.O.A.C.  
Use dilution confirmation.....1:256

**Virucidal** against certain lypophylic viruses including Herpes simplex, Influenza A2, canine distemper, fowl laryngo-tracheitis and Vaccinia, also virucidal against Adenovirus Type 5.  
Use dilution confirmation.....1:256

**DIRECTIONS FOR USE**

	Use Dilution	PROQUAT	Water
General cleaning and disinfection	1:256 (470 ppm)	20 mL 2L	5.0 L 500 L

**Poultry and Livestock Buildings, Vehicles and Equipment:**

Remove all poultry and livestock from buildings. Remove all litter and manure from floors, walls and other surfaces of barns and equipment. Apply PROQUAT at the recommended use dilution and saturate all surfaces by spray or manual methods. Preliminary cleaning may be necessary. Ventilate buildings and allow disinfectant to dry before housing livestock. Rinse feeders and waterers thoroughly with potable water prior to reuse.

**Hatchery Sanitation:**

Use PROQUAT 1:256 to clean and disinfect hatchers, equipment, floors and walls. Also for disinfection of setter and hatcher trays before reuse. Saturate all surfaces. Preliminary cleaning may be necessary.

**DISPOSAL**

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Environment Canada.
5. For small spills. Recover free liquid. Soak up residue with

solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN #4.

**NOTICE TO USER:**

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN.

DANGER - corrosive to eyes. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed.

Avoid contamination of food.

**FIRST AID INSTRUCTIONS:**

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, or gelatin solution. If these are not available, drink large quantities of water. Avoid alcohol. Never give anything by mouth to an unconscious person. Call a physician immediately.

**TOXICOLOGICAL INFORMATION:** Probable mucosal damage may contraindicate inducing vomiting. Measures against circulatory shock, as well as oxygen and measures to support breathing manually or mechanically may be needed. If persistent, convulsions may be controlled by cautious intravenous injection of a short-acting barbiturate drug. Oral LD50 (rat): 2100 mg/kg.

\* Trademark: **PROQUAT**

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

