

1997

Net Contents 4.5 L

CCS 06-3569-01-000

PROSOVET(R)

GERMICIDAL DETERGENT

GERMICIDAL, FUNGICIDAL, VIRUCIDAL

Recommended for the cleaning and disinfecting of livestock and poultry buildings, vehicles and equipment, and hatcheries and equipment. Do not use on food contact surfaces.

GUARANTEE:

o-Phenylphenol 9.20%,  
o-Benzyl-p-chlorophenol 8.20%  
p-Tert Amylphenol 1.70%

AGRICULTURAL

REGISTRATION NO. 15439 PEST CONTROL PRODUCTS ACT

READ THE LABEL BEFORE USING

DANGER

CORROSIVE POISON

Pharmacia & Upjohn Animal Health  
40 Centennial Road  
Orangeville, Ontario L9W 3T3

PROSOVET Concentrate Germicidal Detergent is a highly concentrated disinfectant-cleaner with broad spectrum antimicrobial activity effective in soft or hard water (tested at 400 ppm CaCO<sub>3</sub>). No need to add soap or other detergents.

Germicidal Staphylocidal Pseudomonacidal Tuberculocidal A.O.A.C.  
Use dilution confirmation ..... 1:256  
Fungicidal T. interdigitale A.O.A.C.  
Use dilution confirmation ..... 1:256  
Virucidal against certain lipophilic viruses including influenza

A2, Herpes simplex, Vaccinia and the causative agent of Newcastle Disease. It is also virucidal against Adenovirus Type 2.  
 Use dilution confirmation ..... 1:256

DIRECTIONS FOR USE:

	Use Dilution	PROSOVET Concentrate	Water
General cleaning and disinfection	1:256 (0.39%)	40 mL 2 L	10 L 500 L
Heavily soiled areas	3:200 (1.5%)	150 mL	10 L

Avoid waste: dilute as directed.

Poultry and Livestock Buildings and Equipment, Vehicles:

Remove all animals from premises. Remove all litter and manure from floors, walls and other surfaces of barns, vehicles and equipment. Empty all feeding and watering equipment. Apply PROSOVET Concentrate at the recommended use dilution (1:256) and saturate all surfaces by spray or manual methods. Ventilate buildings and allow disinfectants to dry before housing livestock. Rinse feeders and waterers thoroughly with potable water prior to reuse. In heavily soiled areas use the 3:200 dilution of PROSOVET Concentrate.

Hatchery Sanitation: Remove all poultry from premises. Remove all litter and droppings from floors and surfaces. Use PROSOVET Concentrate diluted 1:256 to clean and disinfect hatchers, equipment, floors and walls. Also for disinfection of setter and hatcher trays before reuse. Saturate all surfaces. In heavily soiled areas increase solution strength to 1.5%.

Foot Baths: Fill entry baths with fresh solution of PROSOVET Concentrate 3:200 dilution to a depth of about 3 cm. Remove excess soil with broom or scraper before entering foot bath. Change solution when noticeably dirty.

Disposal:

1. Rinse the emptied container thoroughly and add the rinsings to treatment site.
2. Follow provincial instructions for any required additional

- cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
  4. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Environment Canada.
  5. For small spills. Recover free liquid. Soak up residue with solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN #4.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN.

DANGER - corrosive to eyes. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. Do not breathe vapour fumes.

FIRST AID INSTRUCTIONS: In case of contact immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, or gelatin solution. If these are not available drink large quantities of water. Never give liquids by mouth to an unconscious person. Call a physician immediately.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate inducing vomiting. Measures against circulatory shock, respiratory depression and convulsion may be needed. Acute oral LD50 (rat): 3.65 g/kg.

\* Trademark: PROSOVET

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this

service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))