

01-FEB-1999

AGRICULTURAL

750 g

AMBUSH(R) 25WP

Wettable Powder Insecticide

For Apples, Pears, Peaches, Nectarines and Grapes

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

GUARANTEE: Permethrin 25%

REGISTRATION NUMBER:14964

PEST CONTROL PRODUCTS ACT

ZENECA Agro, a business of ZENECA Corp.  
250-3115 12<sup>th</sup> Street N.E.  
Calgary, AB  
T2E 7J2

A93/A-1

#### PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes and skin. Apply using ground equipment only. Do not apply by aircraft. This product is very toxic to fish and aquatic organisms. Do not contaminate ponds, lakes, streams or rivers during sprayer filling or rinsing operations or while spraying. Do not apply within 15 m of productive fisheries water or water fowl habitat.

Do not apply when weather conditions favour drift from the target area. This product is very toxic to bees, avoid spraying when bees are foraging.

#### FIRST AID

If in contact with eyes, wash with plenty of clean water for at

least 15 minutes and see physician. If on skin, wash with soap and water. If swallowed, induce vomiting and see physician.

#### EMERGENCY TELEPHONE NUMBER

All hours, 1-800-327-8633 (FASTMED) ONLY for health and environmental information.

#### DISPOSAL:

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

#### PRODUCT INFORMATION

AMBUSH 25WP is a synthetic pyrethroid insecticide of low mammalian toxicity. It is a fast acting stomach and contact insecticide with no systemic or fumigant effect. It is recommended for control of insects as specified below on apples, pears, peaches, nectarines, and grapes. For all crops listed below, use sufficient water for thorough coverage. Follow provincial spray calendars for optimum timing of spray applications. Where a rate range is suggested, use the higher rate only when heavy insect pressure is anticipated. NOTE: REPEATED APPLICATIONS ARE NOT ADVISED IN ORCHARDS WHERE INTEGRATED PEST MANAGEMENT IS IN EFFECT BECAUSE SEVERE REDUCTIONS OF BENEFICIAL ARTHROPODS MAY RESULT.

#### NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

#### DIRECTIONS

GROUND APPLICATION ONLY

APPLES: Winter Moth, Eastern Tent Caterpillar - Apply 425 g/ha of AMBUSH 25WP. Eye-spotted Budmoth: Apply 425 to 850 g/ha of AMBUSH 25WP. Plum Curculio, White Apple Leafhopper - Apply 550 to 850 g/ha of AMBUSH 25WP. Tentiform Leafminer, Green Fruitworm, Apple Maggot, Leafrollers, Tarnished Plant Bug, Mullein Plant Bug, Lesser Appleworm, Codling Moth: Apply 850 g/ha of AMBUSH 25WP. Follow provincial spray calendars for optimum timing of spray applications. Do not apply less than 7 days before harvest.

BLUEBERRIES (Vegetative year only): - Thrips - apply 280 g/ha in one application between mid May and early June when plants are 1-2 cm tall.

GRAPES: Grape Berry Moth - Apply 550 g/ha of AMBUSH 25WP. Grape Leafhopper - Apply 275 g/ha of AMBUSH 25WP. Do not apply less than 21 days before harvest.

PEARS: Eastern Canada - Pear Psylla (adults and nymphs): Codling Moth, Green Fruitworm - Apply 850 g/ha of AMBUSH 25WP. Do not apply less than 7 days before harvest. British Columbia - Pre-bloom Pear Psylla (overwintering adults): Apply 850 g/ha of AMBUSH 25WP up to 1 cm green stage of pear foliage growth. Do not use in British Columbia for control of nymphal stages of pear psylla. Post-Bloom - Pear Psylla: Apply 850 to 1700 g/ha of AMBUSH 25WP. Do not apply less than 7 days before harvest at the 850 g rate and less than 14 days before harvest at the 1700 g rate. Codling Moth: Apply 850 g/ha of AMBUSH 25WP. Do not apply less than 7 days before harvest. PEACHES and NECTARINES: Oriental Fruit Moth - Apply 425 to 850 g/ha of AMBUSH 25WP. Plum Curculio and Plant Bugs - Apply 850 g/ha of AMBUSH 25WP. Do not apply less than 7 days before harvest.

PLUMS: Plum Curculio - Apply 850 g/ha of AMBUSH 25WP. Do not apply within 7 days of harvest.

COMPATIBILITY: AMBUSH 25WP is compatible with CHARGE, Self-Emulsifying Spray Oil, OMITE(R), Captan, BENLATE(R), DIKAR(R), DIFOLATAN(R), Dodine, PIRIMOR(R), and POLYRAM(R). Other pesticides have not been tested.

#### NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product, and accepts the product on that condition.

AMBUSH is a registered trademark of Zeneca Limited, England.

Canadian Patent No. 1008455

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))