

93.08.17

VAPORETTE

INTEGRAL BUCKLE
FLEA COLLAR FOR DOGS

Kills fleas up to 4 months

DOMESTIC

READ THE LABEL BEFORE USING

REGISTRATION NO: 19333
PEST CONTROL PRODUCTS ACT

GUARANTEE:

Carbaryl.....16%

NET WT.: 14 g

SANDOZ AGRO CANADA INC.
Suite 302, Plaza 4,
2000 Argentia Road
Mississauga, Ontario
L5N 1W1

Vaporette is a Registered Trademark of Sandoz Ltd.

DIRECTIONS

Remove the collar from the package and place around dog's neck. Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Leaving ample room for expansion as the dog grows, cut off any excessive length and discard in garbage. Collar must be worn loosely, but securely enough to prevent easy removal. To release the collar, squeeze buckle between thumb and finger and pull sharply. Collar should be removed when dog is bathed with soap or detergent. Do not use on puppies under 3 months of age. The flea collar starts killing fleas when it is placed around the dog's neck. The fleas on the dog will be killed and new ones

which may temporarily appear on the dog, will also be killed during the 4 months (17 weeks) collar is worn. For continuous protection, replace collar when effectiveness diminishes. Dispose of any excess collar packaging and expired collars with garbage to keep out of reach of children.

PRECAUTION: Do not open inner envelope until ready to use. DO NOT ALLOW CHILDREN TO PLAY WITH ENCLOSED COLLAR OR ENVELOPE. This collar is intended for use as an insecticide generator and is not to be taken internally by man or animal. Not intended for use on humans. Hands should be washed following application. Some animals may be sensitive to this collar. Any collar, when fastened too tightly, may cause skin irritation. If irritation persists after proper adjustment, remove the collar and consult a veterinarian. Do not use any other pesticide on dog collar while collar is worn. Do not use on sick or convalescing animals. Do not use on animals under drug or pesticide treatment. For use on pregnant animals consult a veterinarian.

FIRST AID: In case of poisoning, call a physician immediately. If dust from collar contacts skin, wash thoroughly with soap and water. If dust contacts eyes, flush eyes with water and get medical attention.

TOXICOLOGICAL INFORMATION: Dust in eyes will cause temporary pupillary constriction. Dust release from collar is a cholinesterase inhibitor.

Atropine and homatropine are antidotal.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))