

20-APR-2004

21-Mar-2005 - Notification of label change received. See below for changes.

25-FEB-2008 Notification - Change in Registrant Name

ProTurf®
Granular Fungicide V
Containing Chloroneb

- **Prevents and controls Pythium blight and gray snow mold**

COMMERCIAL

REGISTRATION NO. 11466 PEST CONTROL PRODUCTS ACT

Notification Change

GUARANTEE: Chloroneb 6.25%

WARNING, CONTAINS THE ALLERGEN WALNUT OR THEIR SHELLS

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

Net Contents: 14.86 kg

Notification Change

~~Nu-Gro PR Inc.,~~
10 Craig St.,
Brantford, ON
N3R 7J1

Agrium Advanced Technologies RP Inc.

ProTurf® is a registered trademark used under license from The Scotts® Company

DIRECTIONS FOR USE

RECOMMENDED RATE: 14.86 kg/1,000 m²

DOUBLE RATE: 29 kg/1,000 m²

For use on turfgrass areas

Whenever used on greens and fairways apply to moist foliage in the morning or evening. Sprinkle lightly just before traffic resumes to avoid pickup of material on shoes, golf balls, or maintenance equipment.

To prevent Pythium blight – apply at the RECOMMENDED RATE to moist foliage when high temperatures and humidity favor development. Repeat at 5-7 day intervals as long as these weather conditions persist.

To control Pythium blight – apply at the RECOMMENDED RATE to moist foliage as soon as disease activity is observed; repeat at 5-7 day intervals during high temperature and humidity conditions.

In case of severe disease attack apply at the DOUBLE RATE to moist foliage and continue RECOMMENDED RATE applications at 5-7 day intervals during high temperature and humidity conditions.

On new seedlings, apply at the RECOMMENDED RATE after sowing seed; at first signs of disease apply to moist foliage of newly germinated grass, and repeat at 7-10 day intervals if evidence of disease persists.

When treating to control Pythium blight do not fertilize or water heavily until disease activity is arrested; prolonged contact with foliage enhances effectiveness.

Gray snow mold – apply at the RECOMMENDED RATE to dry or moist foliage prior to first heavy snow (early winter). Repeat in midwinter if turf is free of snow. A final treatment in late winter as the snow melts, extends protection.

In areas where snow coverage is present the entire winter, apply at the DOUBLE RATE prior to the first snow and at the RECOMMENDED RATE when snow melts in the spring.

FIRST AID: If on skin, wash with plenty of water. **If in eyes**, flush with running water for 5-10 minutes and get medical attention or contact a poison control centre IMMEDIATELY. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. May irritate eyes, nose, throat and skin. Avoid breathing dust. Avoid contact with skin, eyes and clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not graze or feed clippings from treated areas to livestock. Do not contaminate water when disposing of equipment washwaters.

DISPOSAL: Thoroughly empty the contents of the container into the application device. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

| Suggested Spreader Settings | | | | |
|---|---------------------|--------------------------|--------------------------|---------------|
| To provide proper distribution, calibrate spreader before application 14.86 kg treats 1,022 m ² at RECOMMENDED RATE (Use 14.86 kg/1,000 m ²) 14.86 kg treats 511 m ² at DOUBLE RATE (Use 29 kg/1,000 m ²) | | | | |
| SPREADER | GROUND SPEED | WIDTH OF COVERAGE | SPREADER SETTINGS | |
| | | | RECOMMENDED | DOUBLE |
| ProTurf [®] Drop | 5 km/h | Overlap wheels | 6 | 8 |
| ProTurf [®] Rotaries R-7, R7X R-8, R8-A (cone 8) | 5 km/h 5 km/h | 2.4 m 2.7 m | G M | K Q |
| Lely – WTR and WRF Models | 7 km/h | 4.6 m | 4 ½ III | 6 III |
| Lely – HR (PTO Model) | 450 rpm @ 7 km/h | 4.9 m | 4 ½ III | 6 III |

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.