CLEAN CROP(R)

METHOXYCHLOR 50W

COMMERCIAL

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING.

REGISTRATION NO. 4878 PEST CONTROL PRODUCTS ACT

GUARANTEE: Methoxychlor 50%

UNITED AGRI PRODUCTS
R.R. #2
Dorchester, Ontario
NOL 1G5

NET: 10 kg

DIRECTIONS FOR USE:

FRUITS: Use 2-4 kg in 1000 L of water for foliage treatment. Do not treat during bloom.

APPLES, APRICOTS, CHERRIES, PEACHES, PEARS, & PLUMS: For control of Cherry fruit flies and Plum curculio, treat after petal fall. Use 2-3 applications at 7-14 day intervals. Follow provincial recommendations for timing.

CURRANTS, GOOSEBERRIES & STRAWBERRIES: For the control of Currant Fruit flies and Strawberry weevils.

VEGETABLES: Apply 1.7-5 kg to 1000 L of water to cover one ha, unless otherwise indicated. Apply when necessary to control pests. The amount required depends on the size of the crop plants. Repeat if needed. Do not feed treated vegetable crop or crop refuse to livestock.

ASPARAGUS: To control Asparagus beetles, use 1.7-4.3 kg in 1000 L of water per ha.

BEANS: To control Bean leaf beetles, Bean weevils, Leafhoppers, and Mexican bean beetles. Do not use on young lima bean plants.

BEETS, CARROTS, EGGPLANT, LETTUCE, PEPPERS, RADISHES, SPINACH & TURNIPS: To control Blister beetles, Flea beetles and Leafhoppers. (May be used on carrots and turnips within one day of harvest if tops are not to be used for food or feed.)

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KALE & KOHLRABI: To control imported Cabbage worms and Flea beetles. Apply at 7-10 day intervals if needed.

CUCUMBERS, MELONS, PUMPKINS & SQUASH: To control Cucumber beetles, Flea beetles, Leafhoppers, and squash bug.

ONIONS: To control Onion thrips. Apply at 7-10 day intervals.

PEAS: To control Pea weevils.

POTATOES: To control Blister beetles, Colorado potato beetles, Flea beetles, and Leafhoppers.

TOMATOES: To control Blister beetles, Fall army worms, Flea beetles, and Tomato fruit worms.

ALFALFA & CLOVER: To control Alfalfa blotch leafminers, Alfalfa weevils, Flea beetles, Grasshoppers, Leafhoppers, and Sweet clover weevils, use 2.25-2.75 kg in 1000 L of water per ha. May be applied during bloom in early morning or late evening when bees are not foraging. Do not apply within 7 days of harvest or grazing.

LIMITATIONS: Do not apply to crops within the pre-harvest intervals specified below.

- 3 days asparagus, cabbage
- 7 days apples, beans, beets, carrots, cauliflower, cherries, onions, peas, pears, plums, potatoes, radishes, turnips
- 14 days broccoli, brussels sprouts, currants, gooseberries, kale, lettuce, spinach, strawberries

21 days - apricots, peaches

BEEF CATTLE, HORSES, SHEEP, GOATS

Do not apply within 7 days of slaughter.

Beef Cattle - horn flies: Treatments are usually effective for about three weeks. Do not apply to dairy cattle. Spray or Dip: Use 0.5 to 1.0 kg per 100L water. As a spray, make sure animals are wetted to the skin. Do not dip calves under 1 month of age. Use lower rate on young animals and calves.

Horses - fleas, mosquitoes, stable fly: Spray 1.0 to 2.0 kg per 100L water. Spray 30 to 60 mL per animal and repeat daily if necessary. Use lower rate on young colts.

Goats, Sheep - lice, ticks (keds): Do not use on young animals. Spray: Use 1.0 to 2.0 kg per 100 mL water. Spray 30 to 60 mL per animal after shearing and repeat at 2 week intervals.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid inhaling spray mist or dust. Use with adequate ventilation. Avoid contact with skin, eyes, and clothing. Wear goggles, chemically resistant gloves, and long sleeved coveralls when handling this product. When using, do not smoke, drink or eat. Avoid contamination of feed and food. Do not contaminate mangers, drinking troughs, or utensils. Wash skin with soap and water after handling or using. Avoid contamination of ponds, lakes, streams, and other bodies of water which contain fish life or which may be used for irrigation or domestic purposes.

FIRST AID: IF ON SKIN, remove contaminated clothing and wash skin thoroughly with soap and water. If irritation develops or persists, contact a physician. IF IN EYES, flush with plenty of water. If irritation develops or persists, contact a physician. IF SWALLOWED, induce vomiting by giving syrup of IPECAC. Repeat until vomit fluid is clear. If IPECAC is not available, give the patient 1-2 glasses of water and cause vomiting by inserting a finger down the throat. Repeat with water until vomit fluid is clear. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Call a physician or poison control center immediately. Take container, label or product name with you to emergency hospital or physician.

TOXICOLOGICAL INFORMATION: Slightly irritating to the skin. Barbituates are antidotal.

DISPOSAL:

- 1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- 2. Make the empty container unsuitable for further use.
- 3. Dispose of the container in accordance with provincial requirements.
- 4. For information on the disposal of unused, unwanted product and the clean-up of spills contact the provincial regulatory agency or the manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes any risk to persons or property arising from the use of handling and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))