

**17-MAR-2004 Notification received of label changes. See below for these changes.
30-JUN-2006 - Notification of Registrant Name Change. See below for details.**

26-MAY-2000

Later's

Sevin(R) 50% W.P.

Wettable Powder

Carbaryl Insecticide. Controls many Insect Pests
on Vegetables, Ornamentals & Lawns

CONTROLS DEW WORMS

DOMESTIC

WARNING POISON

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

REG. NO. 11096 P.C.P. ACT

Guarantee: carbaryl ... 50%

150 g

Notification Change

~~Later Chemicals Ltd.
12080 Horseshoe Way
Richmond, B.C. V7A 4V5~~

Spectrum Brands IP Inc.
~~Nu-Gro IP Inc~~
10 Craig Street
Brantford, ON N3R 7J1

Later's Sevin 50% Wettable Powder

DIRECTIONS:

Later's Sevin(R) 50% Wettable Powder is a concentrated insecticide. When mixed with water it will provide good insect control on Vegetables, Flowers, Shrubs, Trees and Lawns. Spray in the early morning or in the evening.

5 g = 1 tablespoon
1 m = approx. 1 long pace
10 m² = 5 m x 2 m

HOW TO USE:

VEGETABLES:

Mix the mentioned amount of Later's Sevin 50 W.P. in 1 L of water to cover 10 m², unless otherwise specified. Apply when insects or damage appears: give full coverage.

ASPARAGUS: To control Asparagus Beetles on spears. Use 4.5-11 g; on ferns, 9-18 g. Apply full coverage in the post-harvest period. Do not apply more than once every 3 days.

BEAN: To control Mexican Bean Beetles, use 3 g; for Flea Beetles and Leafhoppers, use 6 g; for Climbing Cutworms, use 6.5 - 9 g; for Lygus and Stinkbugs, use 9 - 11 g.

BET, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CARROT, CAULIFLOWER, CELERY, CHINESE CABBAGE, KOHLRABI, LETTUCE, PARSLEY, PARSNIP, RADISH, SPINACH, and TURNIP: To control Cabbage Worms, Flea Beetles, Lygus Bugs, Meadow Spittlebugs, and Stinkbugs, use 4.5 - 9 g.

CORN: To control Corn Borers, Earworms, Fall Armyworms, and Rootworm adults, use 4.5 - 6.5 g; for Climbing Cutworms use 5 g per 10 m of row (10 long paces) in a 25 - 30 cm band.

CUCUMBER, PUMPKIN and SQUASH: To control Climbing Cutworms, Cucumber Beetles, Flea Beetles, Leafhoppers, and Squash Bugs, use 4.5 g. Injury may occur if applied when the leaves are tender or wet.

EGGPLANT, PEPPER, POTATO and TOMATO: To control Colorado Potato Beetles, use 3 g.

PEA: To control Alfalfa Loopers, use 9 - 11 g.

DAYS TO WAIT BEFORE HARVEST:

1 day	a sparagus, c arrot, c orn, e ggplant, p epper, s trawberry, and t omato.
3 days	b eans, b roccoli, Brussels sprouts, c abbage, c auliflower, c elery, c ucumber, h ead lettuce, k ohlrabi, p ea, pumpkin and squash.
7 days	b eet (roots), p otato, r adish and t urnip (roots)
14 days	b eet (tops), Chinese c abbage, l eaf lettuce, p arsley, p arsnip, s pinach and t urnip (tops)

FLOWERS, TREES and SHRUBS:

Spray in the early morning or in the evening. Use 5 - 7.5 g per 1 L of water and apply full coverage. For Scale insects, apply

when crawlers are present. Do not apply on Boston Ivy or Virginia creeper. Injury may occur if applied when the leaves are tender or wet. Controls Bagworms, Birch Leafminers, Black vine weevils, Blister beetles, Boxelder bugs, Cankerworms, Elm leaf aphids, Elm leaf beetles, Flea beetles, Japanese beetles, Lace bugs, Leafhoppers, Leaf rollers, Mealy bugs, Pine sawflies, Plant bugs, Psyllids, Rose aphids, Rose chafers, Rose slugs, Scale insects, Tent caterpillars, Thrips (exposed) and Willow leaf beetles.

TURF and LAWNS:

Use 20 g in 20 L of water to treat 10 m. Controls Ants, Chinch bugs, Climbing cutworms, Earwigs, Fall armyworms, Fleas, Leafhoppers, Millipedes and Sod Webworms. Mow the lawn before spraying. Do not water for 2 days after treatment. Repeat in 2 - 3 weeks if necessary.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Keep children and pets away from treated areas, at least until spray has dried. Do not breathe dust or spray mist. Avoid skin contact. Wash hands before eating, drinking or smoking. Avoid contamination of feed and foodstuffs.

ENVIRONMENTAL HAZARDS:

Do not contaminate streams, ponds and water supplies. Highly toxic to bees. Do not use where bees are foraging. Warn beekeepers to keep bees away from treated areas for one week after treatment. Some damage may occur to tender foliage in the presence of rain or high humidity of several days duration following spraying.

FIRST AID: **If swallowed**, induce vomiting. To do so give 2 glasses of warm water to drink and insert finger or blunt object into mouth and touch back of victim's throat. Contact a doctor or poison control centre at once. Never give anything by mouth to an unconscious person. **IF IN EYES**, flush with water for 15 minutes and get medical attention. **IF ON SKIN**, remove contaminated clothing and wash skin with soap and water. **IF INHALED**, move the patient to a hospital.

TOXICOLOGICAL INFORMATION: Sevin is a moderate reversible cholinesterase inhibitor. Symptoms of poisoning: headache, nausea, vomiting and blurring of vision. A large dose of atropine may be needed to control the symptoms.

STORAGE: Keep product in original container. Store in a cool, dry place away from food.

DISPOSAL: Do not re-use empty container. Wrap container and discard in household garbage.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

SEVIN(R) is a registered trademark of Rhone Poulenc, Nederland, B.V.

INTERNET: www.laters.com

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))