

USE

GROUP 1B INSECTICIDE

CATTLE DUST BAG

REFILL CARTRIDGES

For control of horn flies and lice on cattle and as an aid in reducing face flies.

For use in Cattle Dust Bags.

COMMERCIAL

READ THE LABEL BEFORE USING

REGISTRATION NO. 16772

PEST CONTROL PRODUCTS ACT

GUARANTEE: Coumaphos 1%

NET CONTENTS: 5.67 kg

UNITED AGRI PRODUCTS **CANADA INC.**

789 Donnybrook Drive

Dorchester, Ontario

N0L 1G5

Notification Change

16292

03/04

DIRECTIONS FOR USE

Beef Cattle, Dairy Cattle (Lactating, Non-Lactating):

For control of horn flies and lice on cattle and as an aid in reducing face flies.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, please note that CATTLE DUST BAG Insecticide contains a Group 1B insecticide. Any insect population may contain individuals naturally resistant to CATTLE DUST BAG Insecticide and other Group 1B insecticide. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance:

- Where possible, rotate the use of CATTLE DUST BAG Insecticide or other Group 1B insecticides with different groups that control the same pests.
- Avoid application of more than the indicated number of sprays of CATTLE DUST BAG Insecticide or other insecticides in the same group in a season.
- Use tank mixtures with insecticides from a different group when such use is permitted.
- Insecticide use should be based on an IPM program that includes scouting, record keeping and considers cultural, biological and other chemical control practices.
- Monitor treated pest populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information and to report suspected resistance, contact the Technical Service, United Agri Products, 1-800-265-4624 or at www.uap.ca.

LIVESTOCK SELF-DUSTER:

Suspend duster in areas frequented by cattle or in gateways or lanes through which the animals pass daily. For horn flies, bottom of the bag should hang 10-15 cm below the top line of the cattle. For face flies the bottom of the bag should be at a height to provide optimum dusting of the face. Best results are obtained when cattle are forced to use the bags daily. For lactating dairy cows, the duster should be suspended in a gateway or lane through which the cattle pass when leaving the milking barn.

May be applied up to the day before slaughter. Do not hang duster over feed, mineral, or water troughs.

Do not use inside barns, milking rooms or in the entrance doorway.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Poisonous if swallowed. Harmful if inhaled or absorbed through the skin. Do not breathe dust. Avoid contact with skin, eyes, mucous membranes, and clothing. Wear rubber gloves when handling. Dust in well ventilated areas. Wash thoroughly with soap and warm water after using or handling. Do not contaminate feeds or foods. Do not contaminate any body of water. Wash contaminated clothing with soap and hot water before re-use.

FIRST AID

IF ON SKIN: Remove all contaminated clothing. Wash skin and hair thoroughly with soap and water. Wash clothing before reuse. If irritation persists, get medical attention or contact a poison control centre.

IF IN EYES: Flush with running water for at least 15 minutes while holding eyelids open to help flush out material. Get medical attention or contact a poison control centre.

INGESTION: Drink 1-2 glasses of water and induce vomiting by inserting a finger down the throat. Never induce vomiting or give anything by mouth to an unconscious person. Keep head lower than chest to avoid aspiration into lungs. Call physician immediately or contact a poison control centre.

INHALATION: Remove to fresh air. If breathing is difficult, administer oxygen; if breathing stops administer artificial respiration. Get medical attention immediately.

Take container, label or product name and pest control product registration number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

Active ingredient is a cholinesterase inhibitor. Atropine sulfate is an antidote. 2-PAM is also antidotal and may be administered in conjunction with atropine.

DISPOSAL

Thoroughly empty the contents of the container into the application device. Follow provincial instruction for any required additional cleaning prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.