

94.01.24

Barn and livestock pressurized spray

Insecticide sous pression
pour bâtiments de ferme
et animaux

DISVAP SPRAY

INSECTICIDE
PRESSURIZED SPRAY

INSECTICIDE
SOUS PRESSION

For milkhouses, milking parlors, barns,
beef and dairy cattle.

Pour laiteries, salles de traite, étables,
bovins de boucherie et laitiers.

COMMERCIAL

SHAKE WELL BEFORE USING
BIEN AGITER AVANT USAGE

GUARANTEE

Pyrethrins.....	0.5%
Technical Piperonyl Butoxide.....	1.0%
N-Octyl Bicycloheptene Dicarboximide.....	1.0%

GARANTIE

Pyréthrines.....	0.5%
Butoxide de pipéronyle technique.....	1.0%
N-Octyl bicycloheptène dicarboximide.....	1.0%

READ LABEL BEFORE USING
LIRE L'ETIQUETTE AVANT L'EMPLOI

REGISTRATION NO. 17250
NUMERO D'ENREGISTREMENT 17250

PEST CONTROL PRODUCTS ACT
LOI SUR LES PRODUITS ANTIPARASITAIRES

WARNING
AVERTISSEMENT

FLAMMABLE
INFLAMMABLE

EXPLOSIVE
EXPLOSIF

NET CONTENTS / CONTENU NET 595 g

DISPAR,
DIVISION DE/OF VETOQUINOL CANADA INC.
675 St-Pierre sud,
Joliette, Canada J6E 3Z1

DISVAP SPRAY kills flies before they can breed.

DIRECTIONS: To control house flies, stable flies and gnats.

Space Spray: Kills flying insects in milking parlors, milkhouses and barns. Direct spray upwards to all parts of the room especially windows and other light sources. Apply at a rate of 1 to 2 seconds per 30 m³. Remove all animals from barn before using as a space spray. For best results, close all doors and windows during treatment. Ventilate treated areas thoroughly for at least 30 minutes before animal reentry. Spray only when temperature is under 32°C.

Livestock Spray: To treat beef and dairy cattle against attack of house flies, stable flies and gnats. Spray animals once daily with a fine mist. Spray should be enough to wet ends of the hair only; do not wet the hair or skin. Take care not to spray in animal's eyes. Treat dairy cattle at least 20 minutes before milking. Wash udders before milking. Do not use more often than once a day.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not remain in treated areas. Ensure proper ventilation after sealed areas have

been treated. Do not spray over milk or milking equipment. Cover milk and milking utensils to prevent contamination from spray and dead insects. Any treated surface that may contact food or feed products should be washed thoroughly with potable water after spraying and before reuse. Do not contaminate food, feed or water troughs. Avoid inhalation of spray. Avoid contact with skin and eyes. Wash after using. Product is toxic to fish. Do not contaminate any body of water.

CONTENTS UNDER PRESSURE: Do not puncture or incinerate can. Do not store at temperature above 50'C or in direct sunlight. Do not use in presence of open flame or sparks.

FIRST AID: If sprayed in eyes or on skin flush with clean water. If irritation persists, obtain medical attention. If accidentally swallowed DO NOT induce vomiting and obtain medical attention at once.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe condition.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DISPOSAL: When container is empty, press valve to release all remaining pressure. Dispose of empty container in accordance with provincial requirements.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))