

18-DEC-2006

29-JAN-2008 Notification - Change in Registrant

THROW BAG

**WILSON®
WILSARIN
RAT AND MOUSE BAIT PELLETS
WITH BROMADIOLONE**

RODENTICIDE

DOMESTIC

REG. NO. 18992 P.C.P. ACT

GUARANTEE: BROMADIOLONE 0.005 %

Warning, contains the allergen wheat.

Contains malathion at 0.01% and calcium propionate at 0.3% as preservatives.

WARNING



POISON

READ THE LABEL AND OUTER CARTON BEFORE USING
KEEP OUT OF REACH OF CHILDREN

50 g

Notification Change

~~Spectrum Brands IP Inc.~~,
P.O. Box 21001,
Brantford, ON
N3R 7W9

Sure-Gro IP Inc.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. Before using, see outer carton for full PRECAUTIONS regarding placement, storage and hygiene.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals or pets, may reduce the clotting ability of the blood and haemorrhage may result. **If in eyes,** flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and INDUCE VOMITING by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty containers. Dispose of empty container with household garbage. Dispose of dead rodents by burying or in household garbage. Dispose of unused or spoiled bait in accordance with local requirements.

For additional product information or in case of emergency, spills or fire, call toll free 1-800-268-2806. www.wilsonproducts.ca

Wilson® Registered Trademark of Spectrum Brands Canada, Inc., used under license.

CARTON

WILSON[®]
WILSARIN
RAT & MOUSE BAIT PELLETS
WITH BROMADIOLONE

A single feeding anticoagulant rodenticide.
Controls house mice (*Mus musculus*), Norway rats (*Rattus norvegicus*), roof rats (*Rattus rattus*).
Kills rats resistant to warfarin and pindone baits.
JUST ONE FEEDING KILLS RATS AND MICE!
FOR INDOOR OR OUTDOOR USE.

DOMESTIC

REG. NO. 18992 P.C.P. ACT

GUARANTEE: BROMADIOLONE 0.005%

Warning, contains the allergen wheat.

Contains malathion at 0.01% and calcium propionate at 0.3% as preservatives.

WARNING



POISON

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

8 x 50 g ready to eat bait bags
(1 kg 20 x 50 g ready to eat bait bags)

Notification Change

~~Spectrum Brands IP Inc.,~~
P.O. Box 21001,
Brantford, ON
N3R 7W9

Sure-Gro IP Inc.

REFER TO AND RETAIN THIS CARTON FOR COMPLETE USE AND HANDLING INSTRUCTIONS.

Wilsarin Rat and Mouse Bait Pellets contains a rodenticide that kills mice and rats by reducing the clotting ability of the blood. One day feeding will achieve control if sufficient bait stations are used to supply rodent population.

DIRECTIONS FOR USE: Place feeding stations in locations frequented by rats and mice, near rat holes, runways, burrows or against walls or in corners where mice are seen. Place bait in tamper-resistant bait boxes or protected areas inaccessible at the time of placement to children, pets, domestic animals and wildlife. Replenish bait stations as needed to ensure continuous supply of fresh bait until complete lack of feeding is noted. A shallow container of drinking water encourages consumption of bait. **For rats: Norway rats (*Rattus norvegicus*) and roof rats (*Rattus rattus*):** Place 100 g of bait at 2 to 3 metre intervals. Maintain a constant supply of bait at stations for at least 10 days. **For mice: (*Mus musculus*):** Place 10-15 g of bait at 1 to 2 metre intervals. An uninterrupted supply of bait should be maintained for at least 15 days. One day feeding will achieve control when the baiting treatment utilizes a sufficient number of placements that have been consumed by the rodents. Control will be evident, however, only in the following days.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK. May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS. Keep out of lakes, streams or ponds.

USE LIMITATIONS: Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

FIRST AID: This product contains an anticoagulant chemical which, if taken accidentally by humans, domestic animals or pets, may reduce the clotting ability of the blood and haemorrhage may result. **If in eyes,** flush with water for 15 minutes. **If on skin,** wash with soap and water. **If swallowed,** give patient one or two glasses of water and INDUCE VOMITING by touching back of the throat with a finger or blunt object; repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention. If pet or livestock poisoning is suspected, immediately contact a veterinarian.

TOXICOLOGICAL INFORMATION: Vitamin K₁ in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case, measured by establishing prolonged prothrombin times (P.T.), will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

STORAGE: Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

DISPOSAL: Do not reuse empty containers. Dispose of empty container with household garbage. Dispose of dead rodents by burying or in household garbage. Dispose of unused or spoiled bait in accordance with local requirements.

For additional product information or in case of emergency, spills or fire, call toll free 1-800-268-2806.

www.wilsonproducts.ca

Wilson® Registered Trademark of Spectrum Brands Canada, Inc., used under license.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.