

95.05.31

RODEX PELLETTED BAIT 1

With COV-R-TOX

ENCAPSULATED WARFARIN

KILLS RATS AND MICE

For Control of Rats and  
Mice Indoors and Outdoors

DOMESTIC

READ THE LABEL BEFORE USING

GUARANTEE:

Warfarin            0.025%

REG. NO. 23952

PEST CONTROL PRODUCTS ACT

NET: 453.6 g

HACCO, Inc.  
P.O. Box 7190  
Madison, WI  
53707

#### DIRECTIONS

NORWAY AND ROOF RATS: Use approximately 100-450 grams per bait station in locations frequented by rats or where rats have been known to feed. Baits must be protected from pets, poultry, livestock, etc., by bait boxes or other suitable means. Baits should be examined regularly and replenished before completely eaten. Replace bait if it molds or decomposes.

HOUSE MICE: For controlling mice, use 40-60 grams per bait station and place 3-4 meters apart. Follow the same general directions as for rats, except that the amount of bait placements should be smaller and more placements should be made. Mice are

more difficult to control than rats and complete control may take a longer period of baiting.

**IMPORTANT:** Baiting should be continued as long as evidence of feeding is observed. Assure an uninterrupted supply of fresh bait for a period of not less than 12 days. Where a continuous source of infestation is present from nearby dumps or fields, permanent stations should be used and bait placed in them as often as necessary.

**PRECAUTIONS:** KEEP OUT OF REACH OF CHILDREN, PETS, DOMESTIC ANIMALS and WILDLIFE. Do not expose treated bait in areas accessible to children, dogs, domestic animals, and wildlife except when enclosed within a tamper-resistant bait box. Keep away from feed or food stuffs. Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after handling and before eating or smoking. Keep out of lakes, streams or ponds.

#### FIRST AID

RODEX Pelleted Bait 1 contains as its active ingredient an anticoagulant chemical which, if taken accidentally by humans, domestic animals or pets, may reduce the clotting ability of the blood and internal haemorrhage may result. **If in eyes**, flush with water for 15 minutes. **If on skin**, wash with soap and water. **If swallowed**, give patient 1-2 glasses of water and induce vomiting by touching back of the throat with a finger or blunt object. Repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. Call a Physician or Poison Control Center immediately.

#### TOXICOLOGY

When a human has been known to have accidentally ingested RODEX Pelleted Bait 1, blood transfusions combined with intravenous injections and oral doses of Vitamin K<sub>1</sub> are indicated as in the case of haemorrhage caused by overdoses of Dicumarol. Vitamin K<sub>1</sub> in the form of intravenous, intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**DISPOSAL:** Do not reuse container. Wrap and dispose of with household garbage. **RODENT DISPOSAL:** Dispose of dead rodents in garbage or by burying.

**NOTICE TO BUYER:** Seller's guarantee shall be limited to the terms

set out in the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )