

85.12.31

CIBA-GEIGY

READ THE LABEL BEFORE USING

COMMERCIAL

Contains Rotenone.
For control of Lice, Fleas, Ticks and Mites.

LIVESTOCK LOUSE POWDER

INSECTICIDE

GUARANTEE: Rotenone.....1%

REGISTRATION NO.: 1683
PEST CONTROL PRODUCTS ACT

Net Weight: 1 Kg.

CIBA-GEIGY CANADA LTD.
Agricultural Division
6860 Century Avenue,
Mississauga, Ontario L5N 2W5

DIRECTIONS:

LICE ON BEEF AND DAIRY CATTLE (lactating and non-lactating):
Apply thoroughly to inside of flanks, the escutcheon, base of
tail, lower side of neck, and other regions where lice occur.
Eggs are not affected, so repeat at three week intervals to kill
newly hatched lice.

Note:

1. Do not spray over milk or milking equipment.
2. Do not use in rooms where milk is processed or stored.

SHEEP KEDS AND LICE: Apply powder thoroughly into the wool,
taking care to part the fleece so that dust reaches the skin.

Repeat as necessary.

HOG LICE: Pay particular attention to underside and legs and dust bedding at rate of 250 g/m². Dust hogs lightly at night. Repeat as necessary. Do not treat suckling pigs, nor the sow.

DOGS AND CATS: (Fleas and Lice:) Rub powder well into coat to reach skin. Sprinkle dust in sleeping quarters as well.

LICE ON HORSES: Make a thorough application. A blanket strapped over animal will increase effectiveness of treatment.

POULTRY LICE AND MITES: Dust birds, sprinkle in nests and apply thoroughly in henhouses.

CAUTION: KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash after handling and before eating, drinking or smoking. Do not contaminate food and feed. Do not contaminate any body of water. Toxic to fish. Toxic to hogs if swallowed.

FIRST AID: In case of poisoning, call a physician immediately. If on skin, remove contaminated clothing and wash skin thoroughly with soap and water. If in eyes, flush with water for 5 - 10 minutes and obtain medical attention. If swallowed, give the patient one to two glasses of water, and cause vomiting by giving one dose (15 mL) of syrup of ipecac. If the patient does not vomit within 20 minutes, give a second dose. If syrup of ipecac is not available, give the patient one to two glasses of water and cause vomiting by inserting a finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

DECONTAMINATION AND DISPOSAL: Empty containers thoroughly. Bury in a safe place away from water supplies or deliver them to sanitary landfill dumps in accordance with municipal requirements.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the

terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))