

86.08.26

King
BUG
KILLER
DUST

READ THE LABEL BEFORE USING

300 g

DOMESTIC

GUARANTEE:
5% CARBARYL

FOR CONTROL OF VEGETABLE, FRUIT
AND POULTRY INSECTS

FOR DOGS AND CATS:
TO CONTROL FLEAS AND TICKS

REGISTRATION NO. 10709
PEST CONTROL PRODUCTS ACT

DISTRIBUTED BY

GREEN CROSS
6860 CENTURY AVE.
MISSISSAUGA, ONT.
L5N 2W5
(416) 821-4430

MANUFACTURED BY
KING PESTICIDES LTD.
CAMPBELLVILLE ONTARIO L0P 1B0

DIRECTIONS: For small areas, apply a light even coating of dust to the upper and lower leaf surfaces. For larger areas apply the rate indicated below.

	g Dust per 100 m ²	REMARKS
Potatoes, Tomatoes, Eggplant, Peppers: Colorado Potato Beetle, Potato Flea Beetle, Leaf Hoppers, Fall Army Worm, Tomato Horn Worm, Lygus Bug, Stink Bug	225-450 450	Apply when insects or damage first appears. Repeat in 7-10 days or as necessary. Allow 7 day interval between last dust and harvest.

Cucumbers, Squash, Melons, Pumpkins: Cucumber Beetles, Squash Bugs, Flea Beetles, Leaf Hoppers	300-450	Some leaf injury may occur if treatments are made when young foliage is wet or during prolonged high humidity. Allow 3 day interval between last dust and harvest.
Corn: European Corn Borer Corn Earworm Fall Army Worm	225-375	Make 4 or more application 5 days apart. Start when eggs are laid. Allow 1 day interval between last dust and harvest. Make 4 applications 2-4 days apart. Start at 25% silk showing.
Beans: Mexican Bean Beetles Leaf Hopper Flea Beetle Lygus and Stink Bug	300 350 600	Apply when insects or damage appears. Repeat at 7-10 day intervals. Allow 3 day interval between last dust and harvest.
Strawberries: Meadow Spittle Bug Strawberry Leaf Roller	350-600	Apply when insects or damage appears. Repeat at 7-10 day intervals or as necessary. Allow 1 day between last dust and harvest.
Blueberries: Blueberry Maggot Cherry and Cranberry Fruit Worm Japanese Beetle	225	Apply three weeks before harvest and repeat 10 days later. Allow 1 day interval between last dust and harvest.
Grapes: Grape Berry Moth Leaf Hoppers	450-675	Apply in morning or evening when vines are damp. Allow 3 day interval between last dust and harvest.
In and around Poultry Houses and Farm Buildings: Northern Fowl and Chicken Mites Fleas Bed Bugs	1.25 kg dust per 10 m ²	Repeat as necessary. Do not contaminate nests, eggs, feeding and water troughs.
On Chickens and Domestic Birds: Northern Fowl Chicken Mites Lice Fleas and Fowl Ticks	500 g dust per 100 birds	Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Do not apply within 7 days of slaughter.

PRECAUTIONS: Keep out of the reach of children. Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing. Toxic to bees. Do not apply during pollination. Do not contaminate streams, ponds and water supplies. Dispose empty containers in garbage.

FIRST AID: If swallowed, induce vomiting (repeat until vomit fluid is clear). Bring patient to nearest hospital taking this labelled container with you. In case of skin contact remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention.

TOXICOLOGICAL INFORMATION: Carbaryl is a moderate, reversible cholinesterase inhibitor, atropine is antidotal.

DIRECTIONS: CATS, DOGS, DWELLINGS:

fleas, lice, ticks: Dust entire body beginning at the head and working back. Thoroughly treat areas under legs, around the tail, between the toes and in the ears. Work the dust into the skin. Comb out dead ticks a few hours after treatment. Dust the living quarters. Repeat weekly if necessary.

PRECAUTIONS: Do not use on nursing puppies or on kittens less than 4 weeks of age. Do not continue treatment if skin or nasal irritations develop. Do not treat eyes, face or scrotum.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))