

97.10.29

RHONE-POULENC

A LIQUID FLOWABLE FOR CONTROL OF
DAMPING OFF AND ROOT ROT
CAUSED BY RHIZOCTONIA SOLANI, SEED BORNE
BLACKLEG, SEED BORNE ALTERNARIA BLACK SPOT
AND FLEA BEETLE IN MUSTARD

ROVRAL ST
MUSTARD SEED TREATMENT

CONTAINS IPRDIONE AND LINDANE

AGRICULTURAL

GUARANTEE: IPRDIONE 167 g/L LINDANE 500 g/L

DANGER

POISON

READ THE LABEL BEFORE USING

REGISTRATION NO. 19035
PEST CONTROL PRODUCTS ACT

RHÔNE-POULENC CANADA INC.
2000 ARGENTIA ROAD, PLAZA 3
MISSISSAUGA, ONTARIO L5N 1V9
(905) 821-4450

IN CASE OF SPILLS, POISONING OR FIRE
TELEPHONE EMERGENCY RESPONSE NUMBER
1-800-334-7577 (24 HOURS A DAY).

NET CONTENTS 100 LITRES

ROVRAL is a registered trademark of Rhône-Poulenc Agrochimie, Lyon -
authorized user Rhône-Poulenc Canada Inc.

DIRECTIONS FOR USE:

ROLL DRUM OR STIR WELL BEFORE USING

ROVRAL ST is a liquid dual purpose seed treatment for use on mustard to control fleabeetles, damping off and root rot caused by *Rhizoctonia solani* and "seed borne" Blackleg and *Alternaria* Black spot on emerging seedlings

Rate: 30 mL per kg of seed or 750 mL per 25 kg bag of seed. (The 100 L container treats 3333 kg of seed).

ROVRAL ST treated seed can be mixed with the granular insecticides *Furadan CR-10 and *Counter 5G. Refer to labels of both products for use recommendations and all safety precautions.

METHOD OF APPLICATION:

ROVRAL ST treated seed should not require drying after treatment and can be stored or bagged immediately. Add ROVRAL ST to the seed slowly and mix well for at least 3-5 minutes in treating equipment. Thorough seed coverage is required.

STORAGE CONDITIONS:

Protect from frost.

DISPOSAL:

1. After use, return the container to the point of purchase or designated locations.
2. This container must only be refilled with ROVRAL ST. Do not reuse container for any other purpose.
3. Prior to refilling inspect thoroughly for damage such as cracks, punctures, abrasions, and damaged or worn out threads on closure devices.
4. Check for leaks after refilling and before transportation. Do not refill or transport damaged or leaking containers. If the container is not being refilled, return to point of purchase.
5. For information on the disposal of unused, unwanted product and the clean-up of spills contact the Provincial Regulatory Agency or the Manufacturer.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest

Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN, AVOID CONTACT WITH SKIN AND EYES. Wear chemical resistant gloves and coveralls when handling ROVRAL ST or treated seed. When handling ROVRAL ST and when treating seed or while augering or handling treated seed, work in a well ventilated area and wear a suitable respirator. Wash hands and exposed skin thoroughly after handling the concentrate and after application.

FIRST AID:

Remove person from contaminated area, remove contaminated clothing. Keep patient warm, comfortable, at rest.

Eye Contact: Irrigate with water for 15 minutes. Get medical attention.

Skin Contact: Wash exposed skin with soap and cold water.

Inhalation: If breathing stops, start artificial respiration.

Ingestion: If patient is conscious, wash out mouth. Induce vomiting by touching back of the throat with a finger or blunt object. Do not give milk or oils. Get medical attention.

TOXICOLOGICAL INFORMATION:

Contains Lindane, a chlorinated hydrocarbon which may be rapidly absorbed following oral ingestion or skin contact. Gastric lavage with saline purgative should be employed; oil purgatives are contraindicated. Symptomatic treatment is recommended, including an appropriate anti-convulsant therapeutic (eg. diazepam) if required.

USE RESTRICTIONS:

Do not use treated seed for feed, food or oil processing.

Do not store ROVRAL ST near feed or foodstuffs.

Do not contaminate feed or foodstuffs.

Do not contaminate domestic or irrigation waters.

Treated seed must be labelled as follows:

"This seed has been treated with ROVRAL ST containing iprodione and lindane. Do not use for feed, food or oil processing". The date of treatment must also be included.

Do not store treated seed for more than 6 months. Treated seed stored for periods in excess of 6 months may decrease in germination at a faster rate than untreated seed. Treated seed stored for more than 6 months should be tested for germination before planting.

Thoroughly clean auger after handling treated seed before using same auger for handling commercial or feedgrains.

Leftover treated seed should not be stored where there is a chance of it becoming mixed with untreated seed.

Store treated seed in cool, dry conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))