

1995/2000

((BAG))

MORESTAN(R) 25WP

25% WETTABLE POWDER

MITICIDE, FUNGICIDE

COMMERCIAL

Net Weight 2 Kg

FOR CONTROL OF MITES AND POWDERY MILDEW

GUARANTEE: CHINOMETHIONAT 25%

REGISTRATION NO. 08588
PEST CONTROL PRODUCTS ACT

DANGER - EYE IRRITANT

KEEP OUT OF REACH OF CHILDREN
(See side panel for first aid and danger statements)

FOR COMPLETE DIRECTIONS FOR USE
SEE BROCHURE ON BOTTOM OF BAG

READ THE LABEL AND BROCHURE BEFORE USING

CHEMAGRO LIMITED
256 BRITANNIA ROAD EAST
MISSISSAUGA, ONTARIO
CANADA L4Z 1S6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DISPOSAL:

1. Do not reuse the container.
2. Completely empty the bag into application equipment and add all rinsings to the spray mixture in the tank.
3. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
4. Dispose of the empty bag in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

DANGER

Causes substantial eye injury. Harmful if inhaled. Avoid breathing dust or spray mist. Do not get in eyes or on clothing. WEAR GOGGLES OR FACE SHIELD. Wear latex or Neoprene gloves and rubber apron. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash separately from household laundry before reuse.

Do not contaminate feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: In experimental animals acutely poisoned by MORESTAN, diarrhea and decreased activity were the only symptoms

observed. The patient should be treated symptomatically.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MORESTAN is a Reg. TM of Bayer AG, Germany.

786190

((PAMPHLET))

MORESTAN(R)
25WP

25% WETTABLE POWDER

MITICIDE, FUNGICIDE

COMMERCIAL

FOR CONTROL OF MITES AND POWDERY MILDEW

GUARANTEE: CHINOMETHIONAT 25%

REGISTRATION NO. 8588
PEST CONTROL PRODUCTS ACT

NET WEIGHT 2 KILOGRAMS

DANGER - EYE IRRITANT

KEEP OUT OF REACH OF CHILDREN
(See rear panel for first aid and danger statements)

READ THE LABEL AND BROCHURE BEFORE USING

CHEMAGRO LIMITED
 256 BRITANNIA ROAD EAST
 MISSISSAUGA, ONTARIO
 CANADA L4Z 1S6

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS FOR USE

MORESTAN miticide-fungicide is recommended for residual control of mites, mite eggs and pear psylla on deciduous fruits, non-bearing strawberries, and ornamentals. MORESTAN 25% Wettable Powder mixed readily with water to form a suspension which is usable in all conventional hand or machine-operated sprayers. For best results, apply a thorough coverage before mites or mite eggs are present in large numbers and before mites cause bronzing of the foliage. Apply during good drying conditions. (See Restrictions). Consult your local Agricultural Authorities for detailed information on timing of application for most effective control.

RECOMMENDED APPLICATIONS

CROP	PEST	Dosage MORESTAN 25%WP	REMARKS
BEARING FRUIT Pre-Bloom & Post-Harvest Apples Apricots Cherries Peaches Pears Plums Prune Plums	Mites Mite eggs	625 g to 1.25 kg/1000L	Apply specified dosage per 1000 L of water in early season sprays as required, (delayed dormant, pre-pink or pink). Additional applications may be made to tree foliage after harvest.
	Pear psylla	2 kg/1000 L	

NON-BEARING FRUITS Apples	Powdery mildew Mites Mite eggs	625 g to 1.25 kg/1000L	Apply specified dosage per 1000 L of water as a full coverage spray*. Spray all foliage surfaces including the underside of leaves for complete coverage. Repeat as necessary. For control of powdery mildew, begin
Apricots Cherries Peaches Pears Plums Prune Plums	Mites (except cyclamen mite) Mite eggs	625 g to 1.25 kg/1000L	preventive applications before disease appears and apply at 7 to 14 day intervals. During heavy infection periods, make more frequent applications.
NON-BEARING FRUITS Strawberries	Mites (except cyclamen mite)	1.1 to 1.75 kg/ha	Apply specified dosage per hectare in sufficient water for complete coverage. Repeat as necessary.
ORNAMENTALS Roses Arborvitae Juniper Yew	Mites Mite eggs	625 g to 1.25 kg/1000L	Apply specified dosage per 1000 L of water (1g/L). Spray all foliage surfaces including the underside of leaves for complete coverage. Repeat as necessary.

*This concentration is calculated for conventional hydraulic-type sprayers. When lower volumes of spray are applied per hectare as with low-pressure, low-volume, airplane, or mist-type equipment, increase the concentration of MORESTAN 25% Wettable Powder in the spray mixture in order to apply amount of MORESTAN 25% Wettable Powder per hectare equivalent to a full coverage spray.

RESTRICTIONS

Do Not apply oils after delayed dormant where MORESTAN 25% Wettable Powder is to be used in the pink application.

Do not use spreader-sticker materials with MORESTAN 25% Wettable Powder.

MORESTAN applied within 3 days following a frost or near frost weather conditions and prior to a long wetting period may result in russetting. Cultivars such as MacIntosh, Empire, Spartan, and Idared appear to be more susceptible.

Do not use on other crops used for food and forage. Do not allow sprays to drift onto other crops, including pastures. Do not graze treated areas.

Use only according to label directions.

Under certain adverse weather conditions, particularly those that cause slow drying of spray deposits, injury may occur to foliage of strawberries, and deciduous fruit trees (apricots, apples, cherries, peaches, pears, plums, prune plums) that have new and/or tender growth.

Phytotoxicity has occurred on the foliage and flowers of several varieties of roses, especially during periods of high temperatures. Treat roses only after cutting and before new growth appears. Before treating large plantings for the first time, treat a few plants and observe for 4 to 5 days. Under certain conditions, some injury to other ornamentals may occur.

DISPOSAL:

1. Do not reuse the container.
2. Completely empty the container into application equipment and add all rinsings to the spray mixture in the tank.
3. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
4. Dispose of the empty container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills contact the regional office of Conservation and Protection, Environment Canada.

DANGER

Causes substantial eye injury. Harmful if inhaled. Avoid breathing dust or spray mist. Do not get in eyes or on clothing. WEAR GOGGLES OR FACE SHIELD. Wear latex or Neoprene gloves and rubber apron. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash separately from household laundry before reuse.

Do not contaminate feed or food. Keep out of reach of children.

FIRST AID: In case of poisoning call a physician immediately. Have patient lie down and keep quiet. If swallowed, vomiting should be induced. Administer water freely and induce vomiting by giving one dose (15ml) of syrup of ipecac. If vomiting does not occur within 10 to 20 minutes, administer second dose. If syrup of ipecac is not available, induce vomiting by sticking finger down throat. Repeat until vomit fluid is clear. The patient should be lying down with the head below the foot level and facing down or to one side. Professional medical assistance should be secured immediately. DO NOT INDUCE VOMITING TO AN UNCONSCIOUS PERSON OR TO PERSONS IN A CONVULSIVE STATE. If on skin, remove contaminated clothing and wash skin immediately with soap and warm water. If eyes are contaminated, wash immediately with flowing water for at least 15 minutes.

TOXICOLOGICAL INFORMATION: In experimental animals acutely poisoned by MORESTAN, diarrhea and decreased activity were the only symptoms observed. The patient should be treated symptomatically.

ENVIRONMENTAL HAZARDS

MORESTAN is toxic to fish. Do not apply where runoff is likely to occur. In cleaning of equipment or disposal of wastes, do not contaminate streams, lakes, or ponds with this material.

NOTICE

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

MORESTAN is a Reg. TM of Bayer AG, Germany.

Chemagro Limited
256 Britannia Road East
Mississauga, Ontario
L4Z 1S6

986021

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))