

90.09.28

CLEAN CROP

CAPTAN 5 DUST

COMMERCIAL

READ THE LABEL BEFORE USING

Exposure to Captan may produce long term health effects. To minimize exposure, follow directions outlined on this label.

REG. NO. 5499 PEST CONTROL PRODUCTS ACT

GUARANTEE: CAPTAN 5%

DIRECTIONS FOR USE: To obtain satisfactory results, use good dusting equipment and apply thoroughly to give uniform deposit of dust. May be applied 2 days before harvest, except grapes - 7 days.

VEGETABLES: - CUCUMBERS: To control Anthracnose and scab, use at the rate of 35-65 kg/ha. Apply at the first sign of disease and repeat at intervals of 5-7 days. Use 35-50 kg/ha on young plants and 50-65 kg on mature plants.

PEPPER: To control Anthracnose: Apply as soon as foliage appears. Repeat 7-10 days.

EGGPLANT: To control early blight, phomopsis blight: Apply as soon as foliage appears. Repeat every 7-10 days.

TOMATO (FIELD-GREENHOUSE): To control Anthracnose, early and late blights, gray-leaf spot, Septoria leaf spot: Use 35-60 kg/ha at the first sign of infection and repeat at 7-10 day intervals. For severe infections on mature plants, use 65-85 kg/ha.

BERRIES: Under conditions favorable to fruit rot, continue applications throughout the harvest period.

STRAWBERRY: To control gray-mold, fruit rot, leaf spot, use at the rate of 50-80 kg/ha. Apply when the new growth starts in spring and repeat at weekly intervals.

BLUEBERRY: To control fruit rot and mummy berry, use at the rate of 35 kg per ha. and then at weekly intervals through bloom.

LOGANBERRY: To control cane blight, fruit rot, leaf spot, spur blight, use at the rate of 22-35 kg/ha. Apply when first blossoms open and then weekly intervals through bloom.

GRAPE: To control black rot and downy mildew, use at the rate of 35-50 kg per ha. Apply just before bloom, and in first cover. For downy mildew, an additional application 2-3 weeks later may be needed.

SEED TREATMENT: - POTATO (SEED-PIECE): To control brown eye disease, damping-off, scab and seed decay, use at the rate of 1.5-2 kg per 100 kg of seed by coating seed-piece evenly with a light film of dust. Store treated piece in open crates until dried, if planting is delayed.

CAUTION: When treating, augering or handling seed, work in a well-ventilated area, and wear a respirator, goggles and gloves. Do not contaminate food or feed. Do not contaminate any body of water. Label unplanted seed "Poisonous to man and animals. Do not use for feed or food. This seed has been treated with Captan".

TURF: To control brown patch, damping off, leaf spot, melting out, root rot, use at the rate of 1.5-2.5 kg per 100 m². Apply before infection occurs in early spring. Repeat at 7-14 day intervals throughout the summer.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Do not breathe dust. When handling or applying, wear goggles, respirator, clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling Captan and before drinking, eating or smoking. Shower after use or at the end of the work day. If an accident occurs, remove contaminated clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift.

SYMPTOMS OF POISONING: - FIRST AID: In case of poisoning, call a physician immediately. IF ON SKIN, remove contaminated clothing and wash skin thoroughly with soap and water. IF IN EYES, flush with water for 5-10 minutes and obtain medical attention. IF SWALLOWED, give the patient one or two glasses of water and cause vomiting by inserting a finger down the throat. Repeat with water until vomit fluid is clear. The patient should be lying down with the head below the level of the feet. DO NOT TRY TO CAUSE VOMITING IF THE PATIENT IS UNCONSCIOUS OR IN A CONVULSIVE STATE.

DECONTAMINATION AND DISPOSAL:

1. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
2. Make the empty container unsuitable for further use.

3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted products and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons and property arising from the use or handling of this product and accepts the product on that condition.

UNITED AGRI PRODUCTS
P.O. Box 22116
London, Ontario
N6C 4N0

NET: 10 kg

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))