

87.06.22

LINDANE 25-W

INSECTICIDE - WETTABLE POWDER

AGRICULTURAL

GUARANTEE: GAMMA BHC (from lindane) ..... 25%

REGISTRATION NO. 16209

PEST CONTROL PRODUCTS ACT

DANGER POISON

READ THE LABEL BEFORE USING

GAROX CHEMICALS INC.  
R.R. NO. 1 CALEDONIA, ONTARIO  
N0A 1A0

PRECAUTION: KEEP OUT OF REACH OF CHILDREN. Hazardous if swallowed, inhaled or absorbed through the skin. After handling, wash thoroughly with soap and water before eating or smoking and change to clean clothing. Do not apply or allow to drift to areas occupied by unprotected persons or animals, or to streams, lakes or ponds. Lindane is toxic to fish, birds and other wildlife. Avoid treating areas inhabited by wildlife.

FIRST AID: If swallowed, induce vomiting (finger in throat), repeat until vomit fluid is clear. Rush patient to nearest hospital taking this labelled container with you. In case of skin contact, remove contaminated clothing and wash skin with soap and water. If in eyes, flush with plenty of water for 15 minutes and get medical attention.

TOXICOLOGY: Symptoms of poisoning are : nausea, vomiting, hyperirritability, convulsions and/or coma, and other symptoms typical of organochlorine insecticide poisoning. Barbiturates may be given to control convulsions. Oxygen may be indicated. Keep patient quiet. Avoid use of morphine and adrenalin.

DISPOSAL: Burn empty bags; stay out of smoke.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

Seller's guarantee shall be limited to the terms set out on the label and subject thereto. The buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

#### DIRECTIONS FOR USE

BEEF CATTLE, HORSES, SHEEP (for PIGS see below): For lice and ticks mix 250-375 g in 200 litres of water and apply as overall spray at 4.5-9 litres per beef cow and 2.25-4.5 litres per horse or sheep; for lice repeat 14-18 days later. The same spray may be used for hornflies on beef only by spraying back and shoulders at 1-2 litres per cow.

For dip-treatment of sheep mix 625 g in 1000 litres of water.

For lice and mangemites on PIGS use 410 g per 200 litres of water and apply as an overall spray at 1-2 litres per animal; for lice repeat 14-18 days later.

DO NOT USE: ON CALVES OR COLTS UNDER 3 MONTHS OF AGE: ON SICK OR WEAK ANIMALS: ON DAIRY ANIMALS. DO NOT USE IN MILK ROOMS. DO NOT USE ON MEAT ANIMALS WITHIN 30 DAYS OF SLAUGHTER. DO NOT USE AS DIP-TREATMENT OF SHEEP WITHIN 60 DAYS OF SLAUGHTER.

TOBACCO: For wireworms apply 35 h in 200 litres of transplant water and use 2500 litres per hectare.

GREENHOUSE ORNAMENTALS AND FLOWERS: For aphids, thrips and white flies spray plants and soil at first sign of damage. Use 30-50 g per 200 square metres in sufficient water for good coverage.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))