

85.12.31

SANEX

malathion 25WP

WETTABLE POWDER INSECTICIDE

For the Control of Aphids, Leafhoppers
and Certain other insects on Fruit,
Vegetables and Ornamentals.

CAUTION POISON

AGRICULTURAL

NET CONTENTS

READ THE LABEL BEFORE USING

REGISTRATION 17451
PEST CONTROL PRODUCTS ACT

GUARANTEE:
Malathion 25%

Manufactured by
SANEX INC.
6439 NETHERHART RD.,
MISSISSAUGA, ONTARIO L5T 1C3

Branches Throughout North America

PRECAUTIONS:

Keep out of reach of children. Hazardous if swallowed. Avoid inhalation of dust or spray mist. Wash exposed skin with soap and water after spraying. Remove contaminated clothing and clean before re-use. Use a proper respirator when spraying in greenhouse or other enclosed area. Do not contaminate feed or foodstuffs. Do not store near food or feed. Highly toxic to fish and bees. Do not contaminate water supplies. Do not treat any plant while in bloom. Partly used bag should be kept tightly sealed when not being used.

FIRST AID INSTRUCTIONS:

If swallowed, cause vomiting; get medical attention. If in eyes, flush with

water; get medical attention. If on skin, wash with soap and water.

Do not try to cause vomiting if the patient is unconscious or in a convulsive state.

TOXICOLOGICAL INFORMATION:

Malathion, upon repeated or prolonged use, may cause cholinesterase inhibition. Atropine is antidotal.

Do not use 2-PAM.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This package is sold by weight, not by volume. Packed as full as practicable by modern automatic equipment, it contains full net weight indicated. If it does not appear full when opened, it is because contents have settled during shipping and handling.

DIRECTIONS FOR USE:

NOTES:

1. Because of variations in growing conditions and insect species across Canada, local spray guideline or agricultural authorities should be consulted for best use of the product in your area.
2. Higher rates are recommended for heavy infestations.
3. Thorough spray coverage of crop foliage is essential; unless otherwise indicated, use sufficient water for thorough coverage.

| CROPS | INSECTS | RATE/HA | INTERVAL BETWEEN LAST SPRAY HARVEST |
|-------|---------|---------|---|
| | | | |

| | | | |
|---|---|----------------------------------|--------|
| Apples, Pears | Aphids Eye-spotted bud-moth, Mealy bug | 6.75 to 13.5 kg | 3 days |
| | Plant bugs | 3.5 to 6.75 kg | |
| Pears | Pear psylla (suppression only) | 6.75 to 13.5 kg | |
| Strawberries | Leafroller Aphids Strawberry Root Weevil (Adults) | 4.5 kg | 3 days |
| Apply in 2250 L water per hectare | | | |
| Currants, Gooseberries | Currant fruit worm (maggot) | 2.5 kg per 1000 L of water | 3 days |
| Apricots, Peaches, Lecanium scale Grapes (in B.C.) | | 9.75 kg | 3 days |
| Asparagus | Asparagus beetle | 5.5 kg | 1 day |
| Beans | Leafhopper, Mexican bean beetle, Aphids | 4.5 kg | 3 days |
| Beets | Leafminer, Springtail | 4.5 kg | 7 days |

| | | | |
|---|--|-------------------|---------------------------|
| | | | |
| Celery | Leafhopper, Aphids | 4.5 kg | 7 days |
| | | | |
| Cucurbits | Aphids, Leafhopper, Cucumber beetle | 4.5 kg | 3 days |
| Do not apply when foliage is wet | | | |
| | | | |
| Field Lettuce (leaf and head) | Aphids, Leafhoppers | 3.5 to 4.5 kg | head - 7 d leaf - 14 d |
| | | | |
| Greenhouse Lettuce Aphids, Thrips, Mites (leaf and head) | | 3.5 to 4.5 kg | head - 7 d leaf - 21 d |
| Use not more than two applications | | | |
| | | | |
| Pepper | Pepper weevil (maggot) | 5.5 kg | 3 days |
| | | | |
| Spinach | Leafminer, Aphids | 5.5 to 6.75 kg | 7 days |
| | | | |
| Tomato | Aphids | 4.5 kg | 3 days |
| | | | |
| Onion | Thrips | 4.5 kg | 3 days |
| | | | |
| Cabbage, Eggplant, Aphids Pea, Potato | | 3.5 to 4.5 kg | 3 days |

| | | | |
|---|--|--|--------|
| Pea - Do not spray when foliage is wet | | | |
| Broccoli, Brussel's Aphids Sprouts, Kale, Radish, Tobacco | | 3.5 to 4.5 kg | 7 days |
| Ornamental | Aphids, Mites, Leafhoppers, Plant bugs, Tent cater- pillars, European pine shoot moth, Whitefly, Bagworms | 3.5 to 5.5 kg | None |
| Mushroom Houses | Phorid and Sciarid flies | 4 to 6 kg per 1000 L of water | None |
| Apply to wooden surfaces to point of run-off as a spray or paint. Avoid contaminating mushroom belts when time of application differs from that recommended for bed applications. | | | |
| Mushroom Beds | Mites, Phorid and Sciarid Flies | 4 to 6 kg per 1000 L of water | 2 days |
| Apply 12 L spray mix per 100 m2 of bed surface immediately after pickling. Repeat approximately twice a week or as required. | | | |

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA

does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))