## SANEX(R)

#### DOMESTIC

## FLY KILLER INSECTICIDE

Kills flies, mosquitoes, gnats, and certain other small flying insects.

NET CONTENTS: 180 g

## **GUARANTEE:**

Pyrethrin							0.18%
Technical piperonyl butoxide							10.0%
D-trans allethrin							1.62%

#### READ THE LABEL BEFORE USING

REG. NO.: 22,153 PCP ACT

DANGER
EXTREMELY FLAMMABLE
EXPLOSIVE

SANEX AGRO INC. 36 HEAD STREET DUNDAS, ONTARIO L9H 3H3

DIRECTIONS: When used as directed SANEX FLY KILLER INSECTICIDE controls flies, mosquitoes, wasps, silverfish, cockroaches, fleas, ants, spiders, crickets, bed bugs, gnats, adult moths, and accessible stages of stored product insects.

Flies, mosquitoes, wasps, hornets, and ants: Close doors and windows. Direct mist upwards to all parts of the room, especially windows and other light sources. Commence application with 3 to 5 bursts(3 to 5 seconds) for every 30 cubic meters of

space, keeping doors and windows closed for 15 minutes after the treatment. Also direct mist into areas where insects are found.

To control flying moths: Treat all enclosures arranging clothing and other articles so that the mist can circulate freely. Apply for 3 seconds per 30 cubic meters of space. Close treated areas for at least 45 minutes.

To control bed bugs: Spray mattress lightly, particularly around seams and tufts. Take beds apart and spray the frames, springs and all joints. Also spray cracks, crevices, baseboards, and other possible hiding places around the room. Repeat as required.

To control fleas & brown dog ticks: Spray infested areas such as pet's bed and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Remove old bedding and replace with fresh clean bedding after treatment. Insects must be contacted to be killed. Do not treat animals with this product.

If Product is Not for Use on Pets (i.e., Premise Treatment for Fleas and Ticks)

- Do not treat pets with this product. For effective flea and tick control, pets should be treated with a registered flea and tick control product designed for use on pets (N.B. this product is not for use on pets), and treatment of the pet must be combined with the cleanup and sanitation of any area used by the pet. Vacuum floors, carpets, and furniture (discard vacuum bag after use) and wash the pet's bedding, living quarters, and surrounding areas prior to treatment with this product.

### PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. Avoid contact with skin, eyes, and clothing. Avoid inhalation of spray. Wash after use. Cover fish bowls and remove pets for one hour when spraying rooms. Remove or cover all exposed foods and cooking utensils before spraying. This dispenser contains a liquid under pressure. Do not puncture or incinerate, or expose the container to temperatures higher than 50 degrees Celsius. Do not spray closer than 1 metre to fabric to avoid staining.

# FIRST AID:

If on skin, wash thoroughly with soap and water. If in eyes, flush with potable water for 15 minutes. If irritation persists, obtain medical attention. Product contains petroleum distillate. If swallowed, DO NOT induce vomiting. Obtain medical attention at once.

## TOXICOLOGICAL INFORMATION:

Treatment for the various reactions is symptomatic.

## DISPOSAL:

When container is empty, press valve to release all remaining pressure. Dispose of empty container in household garbage. (R)Registered trademark of Sanex Agro Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))