

99/2001

Johnson (R)

BOLT(R)

Commercial Rodenticide

for rats and mice

REG. NO. 15079 P.C.P. ACT

85 g

GUARANTEE:

Warfarin 3-(a-Acetylbenzyl)-4-hydroxycoumarin0.025%
N-(2-Quinoxaliny) sulfanilamide (sulfaquinoxaline)....0.025%

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

READ LABEL BEFORE USING.

Manufactured For
S.C. JOHNSON AND SON, LIMITED,
BRANTFORD, ONT., CANADA N3T 5R1

(R) Can. Trade Mark Reg'd
449 432-1278

Bolt Commercial Rodenticide has been field tested in areas densely infested with rats, and which offered a liberal selection of foods.

Bolt Commercial Rodenticide is

Attractive - Field tests have shown that Bolt Commercial Rodenticide is highly acceptable.

Effective - Field tests have shown that Bolt Commercial Rodenticide will kill rats and mice when used as directed.

Results - Long Term Control by continuous baiting with Bolt

Commercial Rodenticide.

CAUTION: Protect children, domestic animals and wildlife from bait. If swallowed by humans, domestic animals or wildlife this material may reduce the clotting ability of the blood and cause bleeding in such cases intravenous and oral administration of Vitamin K combined with blood transfusions are indicated as in the case of hemorrhage cause by overdoses of Bishydroxycoumarin.

Baits should be placed in areas not accessible to children and pets, wildlife and domestic animals or in tamper proof bait boxes.

Keep away from dogs, pets, other domestic animals and wildlife. Do not reuse container. Destroy when empty.

MADE IN U.S.A.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

Rats: Place two Bolt Commercial Rodenticide throw paks where rats feed, water and/or travel. Placement along walls, in corners, and in concealed locations is desirable. Check bait station daily to assure an uninterrupted supply of bait for not less than 10 days. Continue baiting until all signs of feeding have stopped.

Mice: Place 7 to 14 grams of Bolt Commercial Rodenticide in a shallow, dry container next to burrows in runways or where mice are accustomed to feeding. It is recommended that bait be placed at intervals of 3 to 4 meters. Check bait stations daily to assure an uninterrupted supply of bait for not less than 15 days. Continue baiting until all signs of feeding have stopped. Constant Control. Replace depleted bait. Stale, damp or moldy baits should be replaced with fresh bait promptly. Controls Norway Rat, Roof Rat, and House Mice.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any

guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))