07-JAN-2009 2008-4587 Notification: Change in Registrant Address

## JAGUAR<sup>50</sup> RODENTICIDE

# KILLS RATS & MICE Kills Warfarin Resistant Norway Rats and House Mice

Norway rats and house mice may consume a lethal dose in one feeding with first dead rodents appearing four or five days after treatment begins.

REGISTRATION NO.: 26594 PEST CONTROL PRODUCTS ACT

**COMMERCIAL** 

**WARNING** 



**POISON** 

**GUARANTEE** 

Brodifacoum......0.005%
Contains Denatonium Benzoate
Warning, contains the allergen wheat.
Contains malathion at 0.01%, as a preservative.

### READ THE LABEL BEFORE USING

(see back panel for additional precautionary statements)

#### KEEP OUT OF REACH OF CHILDREN

For indoor use only

Only to be used by certified pest control operators, farmers and persons authorized in government-approved pest control programs.

**Notification Change** 

NET CONTENTS: 5.45 kg (12 lbs.)

3699 Kinsman Blvd., Madison WI 53704 USA

Manufactured by: Motomco Ltd., P.O. Box 8422, Madison, WI 53708 USA

Ph: 608-241-0202

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.

May be harmful or fatal if swallowed or absorbed though the skin. Chemical-resistant gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin

Page 1 of 3 - English

thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and water before reuse. KEEP AWAY FROM FEED AND FOODSTUFFS.

All handlers must wear long-sleeved short and long pants, shoes plus socks, and chemical-resistant gloves when handling this product. Users should remove clothing immediately if pesticide gets inside. Then wash skin thoroughly and put on clean clothing.

**FIRST AID:** If in eyes, flush with water for 15 minutes. If on skin, wash with soap and water. If swallowed, give patient one or two glasses of water and INDUCE VOMITING by touching back of throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If a pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Vitamin  $K_1$  in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**ENVIRONMENTAL HAZARD:** This product is toxic to fish and wildlife. Keep out of lakes, rivers, streams or ponds.

**DIRECTIONS FOR USE:** Use bait to control the Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*) and House mouse (*Mus musculus*) inside of dwellings, farm buildings, food service establishments (non-food areas), granary bins (empty), processing plants (feed and food in non-food areas) and storage areas (non-food). **Food Processing, Food Manufacturing, Food Storage and Food Service Areas:** 

For areas not directly related to food processing: Use only in non-food or non-feed area where feed, food, packaging and handling equipment are never opened or exposed. For areas where feed or food is processed, served, or stored: In meat and food processing plants (processing areas), use only when plant is not in operation. Remove or cover all food, packaging material and utensils before placing bait in baiting stations. Remove all baits and dead rodents before reuse of the plant (processing areas include storage and service).

**USE LIMITATIONS:** Bait MUST either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. DO NOT place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food

To ensure safe use of this product, tamper-resistant bait stations must have the following characteristics:

- \* constructed of high-strength material (e.g., metal or injection moulded plastic) and resistant to destruction by children and non-target animals;
- \* entrance designed so that children and non-target animals cannot reach the bait;
- \* internal structure that prevents bait from being shaken loose;
- \* access panel that fastens securely and locks (e.g. metal screw or padlock);
- \* capable of being securely fastened to a surface (e.g. nailed down); and
- \* clearly labelled: "WARNING POISON".

**SELECTION OF TREATMENT AREAS:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. For indoor use only. Maintain a steady spplu of fresh bait. Remove any spoiled or contaminated bait promptly.

#### **APPLICATION DIRECTIONS:**

**RATS:** Place 100 to 500 grams of bait at intervals of 4.5 to 9 metres per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

**HOUSE MICE:** Place 7 to 14 grams of bait at intervals of 2.5 to 3.5 metres per placement. Larger placements (up to 100 grams) may be needed at points of high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**RATS/MICE:** Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

**DISPOSAL:** Do not reuse empty container. Dispose of unused or spoiled bait in accordance with local requirements. Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**STORAGE:** Store in cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

12906				

\*\*\*\*\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.