

13-JUL-2005 Notification received of a label change. See below for this change.

19-APR-2001

Tomen Agro

COMMERCIAL

Net Contents 20 kg

CAPTAN 10 DUST

FUNGICIDE

READ THE LABEL BEFORE USING

CAUTION: Exposure to Captan may produce long term health effects. To minimize exposure, follow directions outlined on this label.

GUARANTEE: Captan 10%

REGISTRATION NO. 11936 PEST CONTROL PRODUCTS ACT

Arysta LifeScience North America Inc.
~~Arvesta Corporation~~

Notification Change

~~TOMEN AGRO, INC.~~
100 First Street, Suite 1610
San Francisco, CA 94105
tel. #: 415-536-3480

Distributor:
TOMEN CANADA, INC.
Toronto Dominion Center, Suite 3312
Toronto, Ontario M5K 1K7
Canada
tel. #: 416-868-1103

B93/A-2

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Never handle material with bare hands. Use rubber gloves; do not use leather or cloth gloves. Avoid spilling on skin or clothing. Do not breathe dust. Wear a respirator and goggles while mixing, loading and treating seed. Wear long-sleeved and long pants, chemical resistant gloves, goggles and NIOSH approved quarter or half mask cartridge respirator (for particulators) during mixing, loading, application, clean-up and repair activities. When handling or

applying, wear goggles and clothing that completely covers arms and legs to minimize exposure. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling Captan and before drinking, eating or smoking. Shower after use or at the end of the work day. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift. When treating, augering or handling treated seed, work in a well-ventilated area, and wear a suitable respirator, goggles and gloves. Label all unplanted seed potatoes "Poisonous to man and animals. Do not use for food or feed. These potatoes have been treated with Captan."

FIRST AID

If a known exposure occurs or is suspected, immediately initiate the recommended procedures below. If further treatment is required, contact a **Poison Control Centre**, a physician or the nearest hospital.

If Swallowed: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger or blunt object placed on the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious, do not induce vomiting or give anything by mouth.

For Eye Contact: Immediately flush eyes with large amounts of running water for at least 15 minutes. Hold eyelids apart during flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention.

For Skin Contact: Flush all affected areas with plenty of water for 15 minutes. Remove contaminated clothing and shoes. Seek medical attention if skin irritation occurs. Wash clothing before re-use.

If Inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

EMERGENCY TELEPHONE NUMBERS

All hours, 1-800-228-5635 ext. 174 **ONLY** for health and environmental information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams and ponds. Do not apply when weather conditions favour drift of dust

from areas treated. Do not apply directly or indirectly to water. Do not contaminate ponds, lakes, streams, or rivers by cleaning of application equipment or disposal of waste.

STORAGE

Store in a cool, dry place. Protect from excessive heat. Do not contaminate food or feed.

DISPOSAL

Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Thoroughly empty the contents into the application device. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Provincial Regulatory Agency or the Manufacturer. Contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spills and for clean-up of spills.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the **Pest Control Products Act** to use a control product under unsafe conditions.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment or aircraft. Apply at the first sign of infestation and repeat 5 - 10 day intervals as needed to maintain control.

APPLICATION RATES

Apple: Scab, Frog-Eye Leaf Spot- 63 to 84 kilograms per hectare. Consult your provincial spray calendar for timing of applications. Do not apply within 7 days of harvest. Early season sprays may cause injury to foliage of Red Delicious, Winesap and other sensitive varieties.

Cherry (Sour, Sweet): Brown Rot, Leaf Spot (Shot Hole) - 42 to 50.5 kilograms per hectare. Apply to trees in pre-bloom, bloom, petal fall, shuck, first, second and third cover sprays, and immediately after harvest. Leaf injury may result to the Schmidt and Emperor Francis varieties of sweet cherry in post-bloom

sprays. May be applied up to 2 days before harvest of sweet cherries. May be applied up to 5 days before harvest of sour cherries.

Peach: Brown Rot and Scab - 42 to 50.5 kilograms per hectare. Apply in late pink, full bloom, petal fall, shuck fall, cover sprays. Continue applications through harvest if conditions favour brown rot. May be applied up to 2 days before harvest.

Potato Seed Piece Treatment: Brown-Eye Disease, Damping-Off - Use at a rate of 195 grams per 25 kilograms cut seed pieces (780 grams to 100 kilograms of cut seed pieces). Apply in a convenient container or by duster attachment over belt. Coat each seed piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is to be delayed for a day or two, store treated seed pieces in open crates, and provide sufficient ventilation to allow cut surfaces to dry before bagging.

Tomatoes: Anthracnose, Septoria Leaf Spot - Apply 17 to 30 kilograms per hectare at the first sign of infection and repeat at 7 to 10 day intervals. Use 33.5 to 42 kilogram per hectare for severe infestations or on mature plants. May be applied up to 2 days before harvest.

Soil and Greenhouse Bench Treatment: Apply 600 to 975 grams per 100 square metres (84 kilogram per hectare). Work into upper 7.5 to 10 centimetres of soil before planting. For damp-off and root rot diseases of seedling and transplants of roses, other shrubs, flowers, lawnseed beds and vegetables such as beans, peas, celery, crucifers, eggplant, pepper and tomato.

Ornamentals: Rose-Black Spot; Carnation-Leaf Spot; Chrysanthemum-Botrytis Flower Blight, Septoria Leaf Spot - Apply to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

Lawns, Turf (golf courses and sod farms only): Brown Patch, Damping-Off, Leaf Spot, Melting-Out, Root Rot - Use 975 to 1000 grams per 100 square metres. Begin treatments before infection occurs in early spring. Repeat intervals of 7 - 14 days throughout the summer. The maximum application rate to turf (golf courses, sod farms) is 47.5 kgs of Captan 10 Dust per hectare with a maximum seasonal rate of 95 kgs/hectare. DO not apply to home lawns, parks, schools, sports fields and other recreational areas.

Begonia (Tuberous), Daffodil, Dahlia, Gladiolus, Iris (Bulbous) Narcissus, Tulip: Damping-Off, Bulb Rots - Use 180 to 450 grams per 100 metres of row. Treat holes or trenches before planting

and work into the soil to a depth of 7.5 centimetres. Alternatively, use 1.25 to 2.5 kilograms per 100 square metres, and before planting broadcast and work into the soil to a depth of 7.5 centimetres.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the **Pest Control Products Act** to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))