## GERMICIDAL LIFT

Liquid - Disinfectant - Deodorant - Cleaner Staphylocidal - Fungicidal - Pseudomonacidal At Disinfection Rates

## COMMERCIAL

## GUARANTEE

n-Alkyl (60% C14; 30% C16; 5% C12; 5% C18)	
Dimethyl Benzyl Ammonium Chlorides,	0.45%
n-Alkyl (68% C12; 32% C14) Dimethyl Ethyl	
Benzyl Ammonium Chlorides,	0.45%
Tetrasodium Ethylenediamine Tetraacetate,	0.61%
Potassium Hydroxide,	1.02%
Sodium Silicate,	5.88%

Registration No. 15336
Pest Control Products Act

DIN 405477

READ THE LABEL BEFORE USING

CAUTION CORROSIVE

NET CONTENTS LITERS

Les Industries Sanfax(R) Ltee Industries Ltd. 1650 SOUTH SERVICE ROAD, DORVAL, QUEBEC H9P 1H9 (514) 683-7700 TELEX: 05-822566

50 FASKEN DRIVE, SECTION 5, REXDALE, ONTARIO M9W 1K5 (416) 675-1661

MONTREAL - TORONTO - WINNIPEG - EDMONTON - VANCOUVER - MONCTON

A CONSOLIDATED FOODS COMPANY - RESPONSIVE TO CONSUMER NEEDS UNE SOCIETE CONSOLIDATED FOODS - SOUCIEUSE DE SATISFAIRE

AOAC USE DILUTION 12th EDITION, 1975

1.	Staphylococcus aureus	1.16
2.	Salmonella choleraesuis	1.16
3.	Pseudomonas aeruginosa (PRD-10)	1.16

AOAC official fungicidal test, 12th edition, 1975

## DIRECTIONS

For light to moderate cleaning use Germicidal Lift at 12mL per liter of water.

For use in Food Service Areas, Food Processing Plants, Institutional and Industrial Cafeterias, Schools, Hospitals, Nursing Homes, Hotels and Motels use at 60mL per liter of water.

For sanitizing of surfaces use Germicidal Lift at 30 mL per liter of water.

For disinfecting surfaces and equipment against Pseudomonas aeruginosa use Germicidal Lift at 125 mL per liter of water. For Trichophyton interdigitale use Germicidal Lift undiluted.

Before using in food processing areas, food products must be removed from the room or carefully protected. After use, equipment and utensils in the area must be thoroughly rinsed with potable water.

PRECAUTIONS DANGER - EYE IRRITANT KEEP OUT OF REACH OF CHILDREN

FIRST AID: In case of contact, flush eyes or skin with plenty of water. For eyes call physician if irritation persists.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available drink large quantities of water. Call a physician.

NOTICE TO USER: This control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

SELLER'S GUARANTEE shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of handling of this product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label

information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))