

90.12.14

CALO-CLOR(R)

Turf Fungicide

Powder Brand

MERCURIAL TURF FUNGICIDE
SUSPENSION TYPE - WET OR DRY APPLICATION

FOR EFFECTIVE CONTROL OF BROWN PATCH AND SNOW MOLD

GUARANTEE:

MERCUROUS CHLORIDE	60%
MERCURIC CHLORIDE	30%
INERT INGREDIENTS	73%
TOTAL	100%

COMMERCIAL

DANGER POISON

READ THE LABEL BEFORE USING

REGISTRATION NO. 3294
PEST CONTROL PRODUCTS ACT

LOT #

1474

11.33 Kg (25 LB.)

GRACE SIERRA CROP PROTECTION COMPANY
1001 YOSEMITE DRIVE,
MILPITAS, CA 95035

DISTRIBUTED BY: JACK VINCELLI, INC.
5803 AVENUE WESTMINSTER/MONTREAL
H4W 2J9 CANADA

WARNING

POISONOUS IF SWALLOWED,
INHALED OR ABSORBED
THROUGH THE SKIN

FIRST AID

IF ON SKIN: wash thoroughly with soap and water.

IF IN EYES: flush with water for 15 minutes and get medical attention.

IF SWALLOWED: drink milk or egg whites beaten in water or gelatin solution. Repeat until vomit is clear. Take more milk or egg whites. IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably by mouth-to-mouth.

PRECAUTIONS

Keep out of reach of children. Store container tightly closed in a dry ventilated place away from feeds and foodstuffs. Do not store in metal containers, as it is corrosive to metals. Do not use aluminum equipment. Poisonous and corrosive. Wear goggles, respirator, natural rubber gloves and protective clothing when handling. Use with adequate ventilation. May be fatal if swallowed. Causes irreversible eye damage. Do not breathe dust or spray mist. Avoid contact with skin, eyes and clothing. Wash hands thoroughly after handling and before eating, drinking or smoking. Remove contaminated clothing and wash before reuse. Keep children, animals and poultry away from spray area. Do not contaminate neighboring crops, ponds, lakes or streams. Apply when air temperature is below 27°C (80°F). Do not apply in conjunction with soluble fertilizers such as urea, ammonium sulfate, ammonium phosphate or sodium nitrate or with herbicides such as 2,4-D. Do not feed clippings of treated grass to livestock. May kill earthworms. Earthworms killed by mercuric chloride are poisonous to birds.

DISPOSAL PROCEDURES

Follow provincial instructions for any required additional cleaning of the container prior to its disposal.

Make the empty container unsuitable for further use.

Dispose of the container in accordance with provincial requirements.

For information on the disposal of unused, unwanted product and the cleanup of spills, contact the Regional office of Conservation and Protection, Environment Canada.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

DIRECTIONS

SNOW MOLD

For treatment of golf course greens, tees, and approaches only. Preventive treatment: Before ground freezes or before a heavy snow, apply CALO-CLOR to greens and approaches at the rate of 92-153 gm. per 100 sq. m. (3-5 oz./1000 sq. ft.) either in a sand or water carrier. An additional application of CALO-CLOR during the mid-winter thaw assures added protection. Apply uniformly.

DOLLAR SPOT AND BROWN PATCH

PREVENTIVE: Dollar Spot and Brown patch are effectively prevented by treating greens with 31-61 gm. of CALO-CLOR per 100 sq. m. (1-2 oz./1000 sq. ft.) every 7 to 14 days either by sand or water carrier.

CURATIVE: If dollar spot or brown patch is already present, 61-92 gm. of CALO-CLOR per 100 sq. m. (2-3 oz./1000 sq. ft.) is usually sufficient to give control. In periods of extreme heat it is preferable to apply only 31 gm. per 100 sq. m. during the cooler part of the day. If disease still persists, another 31 gm. of CALO-CLOR per 100 sq. m. may be applied within 3 or 4 days. This application should be applied during the evening or cooler part of the day. Apply uniformly.

WET APPLICATION

Determine quantity of CALO-CLOR required, in accordance with dosage recommendations below. Measure this amount of CALO-CLOR into a non-metallic container, add water and stir until a thick uniform slurry is formed. Fill the spray tank, using 20 to 38 liters of water for each 100 sq. m. of turf to be treated. Start the agitator and slowly add the CALO-CLOR slurry. Mix thoroughly before applying. Thoroughly wash sprayer and all metal connections immediately after use.

DRY APPLICATION

Add recommended amount of CALO-CLOR to 10 liters of clean sand for each 100 sq. m. of turf to be treated. Mix thoroughly. Apply evenly to turf so that each 10 liters of CALO-CLOR sand mixture evenly covers 100 sq. m. Wear gloves. Better control may be achieved if a light watering is given after applying the CALO-CLOR sand mixture.

Caution regarding use with fertilizers and weed killers.

Experience has shown that injury attributed to fungicides is often caused by fertilizers or weed killers. Avoid applying CALO-CLOR together with soluble fertilizers such as urea, ammonium sulfate, ammonium phosphate, or with weed killer such as 2,4-D.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))