

2000.07.19

SUPPLEMENTAL LABEL

EMERGENCY REGISTRATION  
FOR DISEASE CONTROL IN CHICKPEAS, DRY PEAS AND  
LENTILS IN MANITOBA AND SASKATCHEWAN  
UNTIL SEPTEMBER 30, 2000

AGRICULTURAL

12.5 KG

BRAVO ULTREX 825DG

AGRICULTURAL FUNGICIDE

WARNING: CAUSES SEVERE EYE DAMAGE

KEEP OUT OF REACH OF CHILDREN

READ LABEL BEFORE USING.

GUARANTEE:

Chlorothalonil (Tetrachloroisophthalonitrile) 82.5%  
Contains 825 grams Chlorothalonil (Tetrachloroisophthalonitrile)  
per kg

REGISTRATION NO. 26581  
PEST CONTROL PRODUCTS ACT

ZENECA Agro,  
a business of ZENECA Corp.  
250-3115 12<sup>th</sup> Street  
Calgary, AB T2E 7J2

PRECAUTIONS

WARNING: Causes Severe Eye Damage. Keep Out of Reach of Children.

DO NOT get in eyes.

Wear coveralls over long pants, and a long-sleeved shirt, goggles or a face shield, apron and chemical resistant gloves during mixing, loading, clean-up and repair activities. Wear pants, a long sleeved shirt and chemical resistant gloves during

application.

DO NOT re-enter treated areas within 48 hours. If required, individuals may re-enter treated areas within 48 hours for short-term tasks not involving hand labour only if at least 4 hours has passed since application, and long pants, long-sleeved shirt, hat and chemical-resistant gloves are worn.

AVOID contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using. DO NOT take internally. AVOID breathing spray mist. This product may produce temporary allergic side effects, characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. DO NOT store near feed or foodstuffs. Store in a cool place. Protect from excessive heat. Apply only to areas specified on label.

#### FIRST AID

In case of eye contact, flush eyes with water for 15 minutes, get medical attention immediately. In case of skin contact, wash exposed areas of skin with soap and warm water. This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. In case of inhalation, remove person to fresh air and get medical attention immediately. Take container, label or product name and Pest Control Product Research Permit Number with you when seeking medical attention.

#### TOXICOLOGICAL INFORMATION

Note to Physician: Persons having an allergic reaction respond to treatment with antihistamines or steroid creams and/or systemic steroids.

#### EMERGENCY TELEPHONE NUMBER

All hours, 1-800-327-8633 (FASTMED) ONLY for health and environmental information.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighbouring areas. DO NOT apply directly to water.

Keep out of lakes, streams, and ponds. DO NOT contaminate water by cleaning of equipment, or disposal of wastes. DO NOT apply

where runoff is likely to occur. DO NOT apply when weather conditions favour drift from areas being treated.

A buffer zone of 100 metres for aerial applications and 15 metres for ground applications to protect surface water bodies including lakes, streams, ponds or sloughs from drift should be observed. For aerial application, use mechanical flaggers only. Provincial buffer zones that are greater than the above buffer zones should be respected.

#### DISPOSAL

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site: 1) Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank. 2) Make the empty, rinsed container unsuitable for further use. If there is no container collection site in your area, dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill and for clean-up of spills.

#### NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

#### PRODUCT INFORMATION

Bravo Ultrex 825DG is an excellent fungicide when used according to label directions for control of a broad spectrum of plant diseases. Bravo Ultrex 825DG can be used effectively in dilute or concentrate sprays. Thorough uniform coverage is essential for disease control.

DO NOT combine Bravo Ultrex 825DG in the spray tank with pesticides, surfactants or fertilizers, unless your prior use has shown the combination physically compatible and non-injurious under your conditions of use.

The required amount of Bravo Ultrex 825DG should be added slowly into the spray tank during filling. With concentrate sprays, premix the required amount of Bravo Ultrex 825DG in a clean container and add to the spray tank as it is being filled. Keep

agitator running when filling spray tank and during spray operations.

#### DIRECTIONS FOR USE

Dosage rates on this label indicate kg of Bravo Ultrex 825DG per hectare, unless otherwise stated. Applications should be made in sufficient water to obtain adequate coverage of foliage. Volume of water to be used will vary with crop and amount of plant growth. Spray volume usually will range from 225-1,600 litres per hectare for dilute sprays and 50-100 litres per hectare for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended unless specific directions are given for a crop.

#### FIELD CROPS

Chickpea: Ascochyta blight - Use 1.8-2.4 kgs per hectare in first application and 1.2-1.8 kgs per hectare in subsequent applications. Make first application at very early flowering and remaining applications at 10 day intervals. DO NOT apply within 48 days of harvest. Apply only with ground application equipment in 220 litres of water per hectare. Do not make more than 3 applications per year.

Dry Peas: Ascochyta blight (*Mycosphaerella pinodes*) - Apply Bravo Ultrex 825DG at a rate of 1.2- 1.8 kgs per hectare, beginning at early flowering. Make a second application at early pod set, around 10 days after the first. Always apply the higher rate when conditions are favourable for disease. If conditions remain favourable for disease, a third application should be made 10 to 14 days later during pod filling. DO NOT apply within 32 days of harvest. DO NOT feed the hay from treated crop to livestock. DO NOT allow grazing of treated crop. Do not make more than 3 applications per year.

Lentils: Ascochyta blight and anthracnose - Apply Bravo Ultrex 825DG at a rate of 1.2-2.4 kgs per hectare beginning at preflowering prior to row closure, make a second application 10 to 14 days after the first application (during bloom period). Do not apply within 48 days of harvest. Do not apply Bravo Ultrex 825DG in combination with POAST (R) herbicide and MERGE (R) surfactant or within 48 hours of the application of POAST and MERGE. Do not feed the straw from treated crop to livestock. Do not allow grazing of treated crop. Do not make more than 2 applications per year.

#### NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the

label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

BRAVO is a registered trademark of a Zeneca Group Company

Poast and Merge are registered trademarks of BASF

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))