RAID

FLEA KILLER PLUS

CAT FOAM

FILLS FLEAS FOR UP TO 8 DAYS

PRESSURIZED SPRAY/DOMESTIC

GUARANTEE:

Pyrethrins......0.15%
Piperonyl Butoxide, Technical......1.30%

REG. NO. 24831 P.C.P. ACT

KEEP OUT OF REACH OF CHILDREN

READ THE LABEL BEFORE USING

CAUTION EXPLOSIVE

140 g

Raid Flea Killer Plus Cat Foam provides rapid flea kill and 8-day residual activity. Effective against cat fleas, dog fleas, American dog ticks and brown ticks. May also be used on dogs sensitive to conventional sprays.

DIRECTIONS FOR USE:

SHAKE WELL BEFORE USING. Start by applying a one-second burst of Raid Flea Killer Plus Cat Foam directly on haircoat near the pet's head, avoiding contact with the eyes. Repeat this procedure making sure to thoroughly moisten, but not soak, the entire haircoat. Use a one-second burst for each half kilogram your cat weighs. Usually 4-5 one-second bursts will cover a 2-3 kg cat. The amount will vary with the size of the pet and density of the haircoat. It is important to apply Raid Flea Killer Plus Cat Foam as evenly as possible over your pet's body because fleas will migrate to untreated areas (e.g. toes, footpads, around ears, etc.). Allow your pet to dry. Do not towel dry as this will remove may of the Raid Flea Killer Plus Cat Foam microcapsules. A blow dryer works well to shorten the drying time. If required, reapply Raid Flea Killer Plus Cat Foam weekly.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid contact with eyes. After using wash hands with soap and water. Do not treat kittens or puppies less than 8 weeks old or nursing females. Consult veterinarian before using on pregnant animals. Do not use on animals under drug or pesticide treatment. Do not use on sick animals. Some animals, particularly cats and kittens, may be sensitive to insecticides. If adverse reactions are observed (e.g. heavy salivation, lethargy, breathing difficulties, or other unusual reactions), bathe pet immediately in warm, soapy water. If symptoms persist, seek medical attention.

FIRST AID: CONTAINS PETROLEUM DISTILLATE. If swallowed, do not induce vomiting, contact a physician or Poison Control Centre immediately. If in EYES, flush with plenty of water. If irritation persists, get medical attention. If on SKIN, wash with soap and water.

STORAGE AND DISPOSAL:

Contents under pressure. Do not puncture or incinerate container. Do not use or store near heat or open flame. Do not store below 0EC (32EF). Exposure to temperatures above 50EC (130EF) may cause bursting. Do not re-use empty container. Press valve to release all pressure. Replace cap and wrap in several layers of newspaper and discard in garbage.

Questions? Comments?
Call 800-558-5566 weekdays 9-9 Eastern Time or write Helen Johnson®

Trade Mark Reg'd S.C. JOHNSON AND SON, INC. Reg'd user S.C. JOHNSON AND SON LIMITED BRANTFORD (ON) Canada N3T 5R1

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))