

21-JUN-2005 Notification received of a label change. See below for this change.

2001/2002

**SERGEANT'S (R)  
DOUBLE PROTECTION**

KILLS FLEAS & TICKS

**SERGEANT'S (R)  
FLEA & TICK COLLAR  
FOR DOGS**

READ THE LABEL BEFORE USING

DOMESTIC

REG. NO. 14928 P.C.P. ACT

GUARANTEE:

Naled	15.0%
Propoxur	2.4%

**DOUBLE PROTECTION**  
NALED provides fast  
VAPOURIZING action!  
Propoxur provides long  
term RESIDUAL action!

RED, Blue, Gold  
Comes in these 3 colors

CONTENTS: 1 COLLAR; 30 g  
FITS NECKS UP TO 60 cm

Notification Change

~~CONAGRA PET PRODUCTS CANADA,  
Cambridge, Ontario  
N1T 1J4~~

**SERGEANT'S PET PRODUCTS INC.**  
**2637 S 158TH PLAZA, SUITE 100** □  
**OMAHA NE 68130** □  
**UNITED STATES**

Made in U.S.A.

**SERGEANT'S (R)  
FLEA & TICK COLLAR FOR DOGS**

## READ THE LABEL BEFORE USING

Kills fleas and non-engorged ticks on dogs and new ones picked up from environment for 4 full months. Collar contains a new sustained release system which combines the insecticidal actions of vapour and powder, working two ways to provide fast action and sustained head to tail protection. Wetting will not impair collar's effectiveness. Use continuously for most effective control.

**DIRECTIONS:** Place collar around dog's neck, adjust for fit, allowing 3 fingers between collar and animal, and buckle in place. The collar must be worn loosely to allow for growth of the dog and to permit the collar to move around the neck. Leave 5-7 cm for extra adjustment. Cut off and dispose of excess length. Wash hands after installing collar on dog. Replace when effectiveness diminishes. Put pouch and expired collar in garbage.

**PRECAUTIONS:** Do not open until ready to use. Collar is intended for external use as an insecticide generator and is not to be taken internally by humans or animals. During storage dust will form on collar. Do not get dust or collar in eyes or mouth. Harmful if swallowed. Wash hands thoroughly with soap and water after handling. Inspect neck area for irritation or adverse reaction every few days when collar is first worn. Some animals may become irritated by any collar if it is applied too tightly. If this occurs, loosen collar. If irritation continues, remove collar and consult your veterinarian. Do not use on sick or convalescing dogs, or dogs under drug treatment. Do not use other cholinesterase inhibitors (other organophosphate or carbamate containing parasiticides) while collar is worn or for a few days after removing collar. Do not use on pregnant animals without first consulting a veterinarian. Do not use on nursing females or puppies under three months of age. Do not use on cats.

### DO NOT ALLOW CHILDREN TO HANDLE COLLAR.

**FIRST AID:** If dust gets in eyes or on skin, flush with plenty of water. If irritation develops or persists or if ingested, obtain medical attention.

**TOXICOLOGICAL INFORMATION:** Propoxur (Sendran (R)) is a cholinesterase inhibitor. Atropine is antidotal.

**DISPOSAL:** Wrap and dispose of any excess collar, packaging and expired collars with garbage to keep out of reach of children.

For cats, use Sergeant's Flea & Tick Collar for Cats.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))