

26-MAR-2007

15-MAR-2007 - Allergen statement update.

25-FEB-2008 Notification - Change in Registrant Name

**WILSON<sup>®</sup>**  
**ROZOL PARA-BLOCK**  
**RAT AND MOUSE BAIT**

RODENTICIDE

DOMESTIC

REG. NO. 18009 P.C.P. ACT

GUARANTEE: Chlorophacinone 0.005%

Warning, contains the allergen wheat.

**Update**

READ THE LABEL BEFORE USING  
KEEP OUT OF REACH OF CHILDREN

450 g

Notification Change

~~Nu-Gro PR Inc.~~  
10 Craig Street  
Brantford, ON  
N3R 7J1  
519-757-0077

Agrium Advanced Technologies RP Inc.

Wilson Rozol Para-Block Rat & Mouse Bait is a weatherproof formulation which can be used in high moisture areas indoors and against the outside walls of buildings.

### **USE LIMITATIONS**

Bait **MUST** either be placed in tamper-resistant bait stations or in locations not accessible to children, pets or livestock. **DO NOT** place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

### **DIRECTIONS:**

**Selection Of Treatment Areas:** Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Maintain a steady supply of fresh bait. Remove any spoiled or contaminated bait promptly.

**Rats:** Norway rat (*Rattus norvegicus*) and Roof rat (*Rattus rattus*) - Divide Wilson Rozol Para-Block Rat & Mouse Bait into sections and place at intervals of 4.5 to 9 meters per placement. Maintain an uninterrupted supply of bait for at least 10 days or until all signs of feeding have stopped. **Mice:** (*Mus musculus*) - Divide Wilson Rozol Para-Block Rat & Mouse Bait into small sections and place at intervals of 3-4 metres per placement. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse feeding have stopped. **Rats/Mice:** Replace contaminated or spoiled bait immediately. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

### **PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN, PETS AND LIVESTOCK.**

May be harmful or fatal if swallowed or absorbed through the skin. Rubber gloves must be worn when handling product and when disposing of dead rodents, unconsumed bait and empty containers. Avoid contact with eyes, skin or clothing. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Wash skin thoroughly with soap and water after handling. Wash contaminated clothing, separately from other laundry, with soap and hot water before reuse. **KEEP AWAY FROM FEED AND FOODSTUFFS.**

This product is toxic to fish and wildlife. The use of anticoagulant formulations can present a significant hazard to predatory mammals and birds through secondary poisoning. Do not contaminate water by cleaning of equipment or disposal of wastes. Keep out of lakes, streams or ponds.

**FIRST AID: If swallowed,** give patient one or two glasses of water and **INDUCE VOMITING** by touching back of throat with a finger or blunt object; repeat until vomit fluid is clear. Do not give anything by mouth to an unconscious person. For all cases of human ingestion, immediately notify a physician or poison control centre. If pet or livestock poisoning is suspected, immediately contact a veterinarian. Take container, label or product name and P.C.P. Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. Vitamin K<sub>1</sub> in the form of intramuscular or subcutaneous injections, or by oral ingestion are suggested remedial

treatments for anticoagulant poisoning. The severity of the case measured by establishing prolonged prothrombin times (P.T.) will determine appropriate therapy. Monitoring P.T. will indicate the necessity of repeated treatments.

**STORAGE:**

Store in a cool, dry place away from other chemicals and food or feed. Store product not in use, in original container, in a secure location inaccessible to children and non-target animals.

**DISPOSAL:** Dispose of empty container in household garbage.

**Rodent Disposal:** Dispose of dead rodents in garbage or by burying.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.