

94.02.17

SANEX

Strychnine 2% Manufacturing Concentrate  
For Use in Formulating Ready-to-Use Baits

DANGER \*\* POISON

READ THE LABEL BEFORE USING

GUARANTEE: 2% Strychnine

REGISTRATION NUMBER 23177 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 4L

MANUFACTURED BY:  
SANEX AGRO INC.  
36 Head Street,  
Dundas, ON L9H 3H3  
(905)-627-9205

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

DIRECTIONS FOR USE: To be used only in the manufacture of rodenticide baits which are registered under the Pest Control Products Act. These baits may only be used for the control of Northern pocket gopher (*Thomomys Talpoides*), Richardson's, Columbian, Franklin and Thirteen-lined ground squirrels (*Spermophilus richardsonii*, *S. columbianus*, *S. franklinii*, *S. tridecemlineatus*, respectively).

PRECAUTIONS: KEEP AWAY FROM CHILDREN and DOMESTIC ANIMALS. Do not contaminate feed or foodstuffs. Do not contaminate water by cleaning of equipment or disposal of wastes. May be fatal if swallowed. Wear protective face mask and rubber gloves.

FIRST AID and TOXICOLOGICAL INFORMATION: OBTAIN MEDICAL ATTENTION IMMEDIATELY. If swallowed, induce vomiting IMMEDIATELY before the onset of convulsions (reflex hyperexcitability).

WARNING: Convulsions may occur within 5 to 10 minutes of ingestion. If more than 5 minutes have elapsed since ingestion, DO NOT induce vomiting. A safer procedure is to ingest a slurry of 30-45 g of activated charcoal in a glassful of water. Gastric lavage is postponed until the patient is fully pre-medicated. DO NOT delay transport of victim to a medical facility. Keep patient in a dark, quiet, warm room with minimal sensory stimulation. Intravenously administer sufficient amounts of barbiturate drugs to prevent further convulsions (amobarbital sodium or pentobarbital sodium). Artificial respiration may become necessary. Keep patient under continual observation until convulsions and reflex irritability have been suppressed and gastric lavage can be performed safely.

DISPOSAL:

1. Rinse the empty container thoroughly and add the rinsings to the formulation tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

LIMITATION OF WARRANTY: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from

reliance upon this service.

+))