

11-APR-2002

# VENTURE™ 25DG

Herbicide  
Dispersible Granules

AGRICULTURAL

Postemergence Herbicide for Control of Grasses in Broadleaf Crops

GUARANTEE:  
fluazifop-P-butyl ..... 25%

**READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

CAUTION



POISON

**POTENTIAL SKIN SENSITIZER**

REGISTRATION NO. **23480**  
PEST CONTROL PRODUCTS ACT.

Net Contents: **6.4 kg**

Syngenta Crop Protection Canada, Inc.  
140 Research Lane  
Research Park, ~~University of Guelph~~  
Guelph, Ontario  
N1G 4Z3

Notification Change

C96/A-4  
**NOTICE TO USER**

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

---

## NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product, and accepts the product on that condition.

## FIRST AID

**IN CASE OF POISONING**, contact a physician or a poison control centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, do not induce vomiting; call a physician immediately or contact a poison control centre.

**If in eyes**, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

**If on skin**, wash with soap and water. If irritation develops and persists, get medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically for ingestion and/or skin and eye contact. Potential skin sensitizer.

## PRECAUTIONS

1. Harmful if swallowed. CAUTION: Potential skin sensitizer. Failure to follow the directions may cause a health hazard.
2. Do not wear contact lenses when using chemicals.
3. All users must follow all handling precautions, use directions, and clean-up procedures noted on the label.
4. Wear coveralls, boots, and PVC (liquid proof) gloves when spraying, or when adjusting, repairing, or cleaning equipment.
5. Wear coveralls, boots, and PVC gloves and safety goggles when handling the granules.
6. Use PVC gloves and dispose of if damaged or after use. Dispose of gloves immediately if contaminated on the inside. Wash hands thoroughly with soap and water before using a new pair of gloves.
7. Avoid spray mist by staying upwind from the spray and/or by wearing a suitable mask or respirator.
8. Wash thoroughly with soap and water after handling and before eating, drinking or smoking.
9. Remove contaminated clothing and wash before re-use.
10. Be sure spray tank is clear of debris and all spray nozzles and screens are clean and clear of obstructions before loading the sprayer with VENTURE 25DG HERBICIDE Herbicide. This will eliminate contamination of clothing and skin by the need to clear blocked nozzles after spraying commences.
11. Avoid drift on to other crops and non-target areas. Corn, cereals and turf are highly susceptible to VENTURE 25DG HERBICIDE Herbicide.
12. Do not apply by aircraft.
13. Do not apply within 15 m of fish-bearing waters and wildlife habitats.
14. Seed only broadleaf crops listed on this label if it is necessary to reseed a crop within 60 days of applying VENTURE 25DG HERBICIDE Herbicide.
15. CAUTION: Unless otherwise noted on the product label, do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

---

## **STORAGE**

Product is not affected by temperatures below 0°C.

## **SPILL CLEANUP**

Follow safety precautions as directed for handling the product. For spillage on floors or hard surface areas, sweep up spilled product for disposal or use. If on floor or hard surface, wash area with detergent and water, and then rinse. Dispose of granules and cleanup materials in an approved landfill site. For information on the disposal of unused, unwanted product contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **DECONTAMINATION AND DISPOSAL:**

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **CONTAINER DISPOSAL:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING  
CALL 1-800-327-8633 (FASTMED)***

---

# VENTURE™ 25DG

Herbicide  
Dispersible Granules

AGRICULTURAL

Postemergence Herbicide for Control of Grasses in Broadleaf Crops

GUARANTEE:  
fluazifop-P-butyl ..... 25%

**READ THE LABEL AND BOOKLET BEFORE USING  
KEEP OUT OF REACH OF CHILDREN**

CAUTION



POISON

**POTENTIAL SKIN SENSITIZER**

REGISTRATION NO. **23480**  
PEST CONTROL PRODUCTS ACT.

Net Contents: **6.4 kg**

Syngenta Crop Protection Canada, Inc.  
140 Research Lane  
Research Park, University of Guelph  
Guelph, Ontario  
N1G 4Z3

C96/A-4

---

## NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

## NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product, and accepts the product on that condition.

## FIRST AID

**IN CASE OF POISONING**, contact a physician or a poison control centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, do not induce vomiting; call a physician immediately or contact a poison control centre.

**If in eyes**, immediately flush with plenty of water for at least 15 minutes. Get medical attention.

**If on skin**, wash with soap and water. If irritation develops and persists, get medical attention.

## TOXICOLOGICAL INFORMATION

Treat symptomatically for ingestion and/or skin and eye contact. Potential skin sensitizer.

## PRECAUTIONS

1. Harmful if swallowed. CAUTION: Potential skin sensitizer. Failure to follow the directions may cause a health hazard.
2. Do not wear contact lenses when using chemicals.
3. All users must follow all handling precautions, use directions, and clean-up procedures noted on the label.
4. Wear coveralls, boots, and PVC (liquid proof) gloves when spraying, or when adjusting, repairing, or cleaning equipment.
5. Wear coveralls, boots, and PVC gloves and safety goggles when handling the granules.
6. Use PVC gloves and dispose of if damaged or after use. Dispose of gloves immediately if contaminated on the inside. Wash hands thoroughly with soap and water before using a new pair of gloves.
7. Avoid spray mist by staying upwind from the spray and/or by wearing a suitable mask or respirator.
8. Wash thoroughly with soap and water after handling and before eating, drinking or smoking.
9. Remove contaminated clothing and wash before re-use.
10. Be sure spray tank is clear of debris and all spray nozzles and screens are clean and clear of obstructions before loading the sprayer with VENTURE 25DG HERBICIDE Herbicide. This will eliminate contamination of clothing and skin by the need to clear blocked nozzles after spraying commences.
11. Avoid drift on to other crops and non-target areas. Corn, cereals and turf are highly susceptible to VENTURE 25DG HERBICIDE Herbicide.
12. Do not apply by aircraft.
13. Do not apply within 15 m of fish-bearing waters and wildlife habitats.
14. Seed only broadleaf crops listed on this label if it is necessary to reseed a crop within 60

- days of applying VENTURE 25DG HERBICIDE Herbicide.
15. CAUTION: Unless otherwise noted on the product label, do not graze the treated crops or cut for hay; sufficient data are not available to support such use.

## STORAGE

Product is not affected by temperatures below 0°C.

## SPILL CLEANUP

Follow safety precautions as directed for handling the product. For spillage on floors or hard surface areas, sweep up spilled product for disposal or use. If on floor or hard surface, wash area with detergent and water, and then rinse. Dispose of granules and cleanup materials in an approved landfill site. For information on the disposal of unused, unwanted product contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## DECONTAMINATION AND DISPOSAL:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## CONTAINER DISPOSAL:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING  
CALL 1-800-327-8633 (FASTMED)***

## PRODUCT INFORMATION

VENTURE 25DG HERBICIDE Herbicide is a selective postemergence herbicide for control of a broad range of annual and perennial grasses in broadleaf crops.

VENTURE 25DG HERBICIDE Herbicide does not control broadleaf weeds, fescue and bluegrass species or sedges including nutsedge.

VENTURE 25DG HERBICIDE Herbicide is a systemic herbicide which is translocated from the treated foliage to the growing points of leaves, shoots, roots or rhizomes. Although the growth of

---

grasses ceases soon after application, the process of killing the foliage, rhizomes and the entire plant may take up to several weeks.

VENTURE 25DG HERBICIDE Herbicide must always be used in conjunction with an adjuvant or poor weed control will result. See ADJUVANTS section. VENTURE 25DG HERBICIDE Herbicide activity is not affected by rainfall if it commences two hours or more after application.

## **DIRECTIONS FOR USE**

VENTURE 25DG HERBICIDE is to be used only on the crops listed on this label.

Do not apply the tank mix of VENTURE 25DG HERBICIDE and MUSTER on Polish varieties of canola as injury may result.

**TIME OF APPLICATION:** VENTURE 25DG HERBICIDE will be less effective when weeds are not growing rapidly because of stress by lack of moisture, excessive moisture, flooding, low temperature and/or very low relative humidity. Regrowth by tillering may occur if application is made under any of the above stress conditions. In wide row crops where the crop canopy may be slow to close, cultivation may be necessary to control grasses that emerge after treatment.

**ANNUAL GRASSES:** Apply VENTURE 25DG HERBICIDE to actively growing annual grasses at the full 2-leaf to 5-leaf stages (except green and yellow foxtail). For green and yellow foxtail, apply at full 2-leaf to 4-leaf stages. Total leaf count should include leaves on the main stem and on tillers; **see illustration 1**. Established grasses beyond the 5-leaf stage (including regrowth after clipping or tilling) will not be controlled. Apply VENTURE 25DG HERBICIDE to green and yellow foxtail before tillering or unacceptable grass control will result. Most effective control of other annual grasses is achieved when application is made before tillering. **Application made to tillered annual grasses that are under moisture and/or temperature stress will not provide acceptable control.**

## **INSERT ILLUSTRATION HERE**

**QUACK GRASS (SUPPRESSION ONLY):** Apply VENTURE 25DG HERBICIDE to actively growing plants that have 3 to 5 fully developed leaves. Application made to plants greater than 20 cm in height or which have reached the heading stage will not provide adequate control. In canola and soybeans only, VENTURE 25DG HERBICIDE applied at lower rates (0.5 kg/ha) will suppress top growth of quack grass throughout the growing season.

**ANNUAL CROPS:** Rhizomes of quack grass should be thoroughly fragmented by tillage (disc or cultivator) prior to the application of VENTURE 25DG HERBICIDE to obtain effective control. Tillage required to fragment rhizomes can be done in the fall or in the spring before seeding. Normal seedbed preparation will fragment quack grass rhizomes sufficiently. Therefore, extra tillage operations are not required. Do not cultivate for 5 days after applying VENTURE 25DG HERBICIDE.

**PERENNIAL CROPS:** In perennial crops, effective seasonal control can be achieved providing that quack grass is not under moisture and/or temperature stress, and that application is confined to the optimum leaf stage. Crop competition generally enhances control of quack grass. Do not cultivate for 5 days after applying VENTURE 25DG HERBICIDE.

## **RATES OF VENTURE 25DG HERBICIDE**

General rates of use for VENTURE 25DG HERBICIDE are given in the "Rates of VENTURE 25DG HERBICIDE" section to control various grass species. Refer to the "Crop Recommendations" section for maximum use rate on each crop, the corresponding spray/harvest interval and recommended tank mixes.

	<b>Rate (kg/ha)</b>	<b>Recommended Leaf Stage</b>
<b>MIXED STANDS of COMMON GRASSES</b>		
<i>Green and Yellow foxtail*</i> , <i>wild oats, wild proso millet,</i> <i>volunteer spring wheat and spring</i> <i>barley, Persian darnel, barnyard grass,</i> <i>and topgrowth suppression of quack grass</i> <i>in soybeans (Eastern Canada) and canola.</i>	<b>WESTERN CANADA</b> 0.7	2-4
	<b>EASTERN CANADA</b> 0.5	3-5

\* In Eastern Canada, if foxtails are present in the mixed stand, use the rate and timing specified for foxtails in the following annual grass section.

#### **VOLUNTEER CEREALS**

<i>Volunteer spring wheat and spring barley</i> Control of other specific grasses under ideal growing conditions	0.4	2-5
---	-----	-----

#### **ANNUAL GRASSES**

Wirestem mulhy in Eastern Canada	1.0	3-5
Wild Oats	0.4	2-5
Wild proso millet	0.5	2-5
Green and Yellow Foxtail in Western Canada	0.7	2-4
Green and Yellow Foxtail in Eastern Canada**	0.5	2-4
Persian darnel and barnyard grass	0.4	2-5
Volunteer corn	0.3	2-5

\*\* (Use 0.7 kg/ha under dry conditions or for heavy foxtail stands)

#### **QUACK GRASS**

Quack grass topgrowth suppression (Canola and soybeans)	0.5	3-5
--	-----	-----

**Note:** An adjuvant is required to be used with VENTURE 25DG HERBICIDE. See the ADJUVANTS section for directions.

**CROP RECOMMENDATIONS:** Refer to the previous list of grass species controlled, found under 'RATES OF VENTURE 25DG HERBICIDE', for the effective VENTURE 25DG HERBICIDE rate. The following chart gives the maximum rate of VENTURE 25DG HERBICIDE allowable for a crop plus other particulars. Remember, in all cases, to apply VENTURE 25DG HERBICIDE at the proper growth stage of the grass and when grasses are actively growing.



<b>CROP RECOMMENDATIONS</b>				
<b>CROP</b>	<b>MAXIMUM VENTURE 25DG Herbicide RATE</b>	<b>MINIMUM INTERVAL TO HARVEST (In Days)</b>	<b>VENTURE 25DG HERBICIDE Herbicide TANK MIXES</b>	<b>REMARKS</b>
Field Crops Canola	0.7 kg/ha	80	1. LONTREL® 2. MUSTER®	Do not apply VENTURE 25 DG later than fifth leaf stage of the crop.  Follow all label directions and precautions for the product you are tank-mixing.  See the section VENTURE 25DG HERBICIDE TANK MIX WITH BROADLEAF HERBICIDES.  CAUTION: Do not apply MUSTER tank mix to Polish varieties of canola as injury may result.
Triazine-tolerant Canola	0.7 kg/ha	80	1. BLADEX® Liquid 2. LONTREL 3. MUSTER	
Creeping Red-Fescue (Seedling & Established)	0.7 kg/ha		ALLY	Fescue grown for seed only. Apply at 2-5 leaf stage of the crop.  Use VENTURE 25DG HERBICIDE at 0.5 kg/ha for seed head control of foxtail barley and smooth brome.
Flax (Including low linolenic acid varieties)	0.7 kg/ha	80		
Flax (Excluding low linolenic acid varieties)	0.2 kg/ha	80	1. Sethoxydim 184g/L	See the restrictions for the tank mix on the following pages.
Flax (Excluding low linolenic acid varieties)	0.2 kg/ha	80	2. Sethoxydim 184 g/L + MCPA Ester 3. Sethoxydim 184 g/L + BUCTRIL M	See the restrictions for the tank mixes on the following pages.
Mustard	0.7 kg/ha	75		Do not apply later than the fifth leaf stage of the crop.
Peas (Field)	0.7 kg/ha	66	PURSUIT®	Crops treated with PURSUIT cannot be grazed or cut for hay. Straw may be feed to lactating dairy animal and other livestock.

<b>CROP RECOMMENDATIONS</b>				
<b>CROP</b>	<b>MAXIMUM VENTURE 25DG Herbicide RATE</b>	<b>MINIMUM INTERVAL TO HARVEST (In Days)</b>	<b>VENTURE 25DG HERBICIDE Herbicide TANK MIXES</b>	<b>REMARKS</b>
Sunflowers	0.7 kg/ha	120		
<b>Legume Forages</b> Alfalfa	0.7 kg/ha	41	May be tank mixed with 2,4-DB	Apply in spring or fall. Treated crop may be fed to lactating dairy animals and other livestock 41 days after treatment.
Red Clover and Birdsfoot Trefoil	0.7 kg/ha	See remarks	May be tank mixed with 2,4-DB	Apply in spring or fall. Do not harvest for feed or graze livestock in the year of treatment. See the section VENTURE 25DG HERBICIDE TANK MIXED WITH BROADLEAF HERBICIDES.
Soybeans (Eastern Canada Only)	*0.7 kg/ha	90	BASAGRAN® FORTE** BASAGRAN FORTE® Plus PINNACLE®** PURSUIT®	Do not apply VENTURE 25DG HERBICIDE later than the third trifoliolate leaf stage of the crop.

\* Use VENTURE 25DG HERBICIDE at 1 kg/ha only if Wirestem Mulhy is present. See the section VENTURE 25DG HERBICIDE TANK MIXED WITH BROADLEAF HERBICIDES.

\*\* WARNING: Up to 10% reduction in control of grass weeds may occur.

**BROADLEAF WEEDS:** VENTURE 25DG HERBICIDE will not control broadleaf weeds. Therefore, a recommended herbicide for control of broadleaf weeds should be applied separately unless a tank mix is otherwise recommended on this label. Do not apply VENTURE 25DG HERBICIDE following a broadleaf postemergent herbicide. Where VENTURE 25DG HERBICIDE is to be used in sequence with a postemergent broadleaf herbicide, other than MUSTER, apply VENTURE 25DG HERBICIDE first and wait at least 3 days before applying the other herbicide. Where MUSTER is to be used in sequence with VENTURE 25DG HERBICIDE, apply MUSTER first, and wait at least 4 days before applying VENTURE 25DG HERBICIDE. Following these sequences will avoid crop injury. Consult the other product labels for application and precautionary statements.

## **VENTURE 25DG HERBICIDE TANK MIXED WITH BROADLEAF HERBICIDES**

---

Read and strictly follow the precautions and use directions on the label of the broadleaf weed herbicide to be used.

Labelled rates of VENTURE 25DG HERBICIDE can be tank mixed with other herbicides for uses as listed on this label (See Crop Recommendations Table and Rates section.). Rates of use and timing of application should be as indicated on each label for products used in the tank mix. For optimum control of grass and broadleaf weeds, the appropriate stages of growth for grass and broadleaf weeds should coincide otherwise control may be lessened. See MIXING INSTRUCTIONS, ADJUVANTS, and SPRAYING INSTRUCTIONS sections for further information.

### **VENTURE 25DG HERBICIDE TANK MIXED WITH OTHER GRASS HERBICIDES**

#### **FOR THE TANK MIXTURES OF VENTURE 25DG HERBICIDE WITH POAST® 184EC OR SELECT® 240EC TO CONTROL WILD OATS, GREEN FOXTAIL, VOLUNTEER SPRING WHEAT AND VOLUNTEER SPRING BARLEY IN CANOLA AND FLAX (WESTERN CANADA ONLY).**

A tank mix combination of VENTURE 25DG HERBICIDE + POAST 184EC or VENTURE 25DG HERBICIDE + SELECT 240EC may be used at lower rates of each product than either product used alone to control wild oats, green foxtail, volunteer spring wheat and volunteer spring barley in canola and flax. Application should be made at the 2 - 5 (total leaves on main shoot plus tillers) leaf stage of the weeds.

#### **Application Rate:**

For the VENTURE 25DG HERBICIDE + POAST 184EC tank mix use 0.20 kg/ha of VENTURE 25DG HERBICIDE + 0.54 L/ha of POAST 184EC. One of the adjuvants, MERGE®, ASSIST® or TURBOCHARGE®, must be added to the spray solution to give a final strength of 0.7% v/v (7 L for each 1000 L of spray solution), or a combination of these adjuvants may be used, provided the final adjuvant concentration adds up to 0.7% v/v. Adjuvant(s) must be added to the spray tank last.

For the VENTURE 25DG HERBICIDE + SELECT 240EC tank mix use 0.20 kg/ha of VENTURE 25DG HERBICIDE + 0.075 kg/ha of SELECT 240EC. One of the adjuvants, AMIGO® or TURBOCHARGE®, must be added to the spray solution to give a final strength of 0.5 - 1.0 % v/v (5 - 10 L for each 1000 L of spray solution), or a combination of these adjuvants may be used, provided the final adjuvant concentration adds up to 0.5 - 1.0 % v/v. Adjuvant(s) must be added to the spray tank last.

Apply in a spray volume of 100 L/ha at a pressure of 275 kPa.

**Flax (Excluding low linolenic acid varieties)- BUCTRIL M or MCPA Ester Tank Mix:** A tank mix of labeled rates of the VENTURE 25DG HERBICIDE + Sethoxydim 184 g/L + BUCTRIL M or MCPA Ester may be used when flax (Excluding low linolenic acid varieties) is growing vigorously and infestations of wild oats, green foxtail, volunteer wheat and volunteer barley are light to moderate. Avoid spraying flax (Excluding low linolenic acid varieties) in hot, humid conditions or crop injury may result. Tank mixes of BUCTRIL M or MCPA Ester may often antagonize the activity of VENTURE 25DG HERBICIDE + Sethoxydim 184 g/L. The following adjuvant or adjuvant combinations, MERGE, ASSIST + MERGE, CHARGE + MERGE, must be added to the spray solution to give a final or final combined strength of 0.7%v/v (7 L for each 1000 L of spray solution). Adjuvants must be added to the spray tank last. Refer to the respective broadleaf weed herbicide label for list of weeds controlled.

---

## VENTURE 25DG HERBICIDE TANK MIX WITH LIBERTY®

THIS TANK MIX MAY ONLY BE USED ON CANOLA VARIETIES, INCLUDING HYBRIDS, THAT ARE TOLERANT TO LIBERTY HERBICIDE. APPLICATIONS TO ANY OTHER CANOLA VARIETIES OR CROPS WILL RESULT IN CROP DEATH.

For control of Volunteer Barley and selected broadleaf weeds apply 0.10 - 0.20 kg of VENTURE® 25DG with 2.67 Litres of LIBERTY® per ha. Add TURBOCHARGE® at 0.5% v/v. Refer to the LIBERTY® label for list of broadleaf weeds and other annual grass weeds controlled and proper application timing.

For further precautionary and application information, refer to the respective product labels.

## VENTURE 25DG HERBICIDE TANK MIX WITH PURSUIT®

To control annual grass and broadleaf weeds in field peas, a tankmix of VENTURE 25DG HERBICIDE + PURSUIT can be used. Use 0.7 kg/ha of VENTURE 25DG HERBICIDE with 0.21 L/ha of PURSUIT in 100 L/ha of water. TURBOCHARGE adjuvant must be added to the spray solution to give a final strength of 0.5 % v/v (0.5 L for each 100 L of spray solution). TURBOCHARGE must be added to the spray tank last. Tank mix of VENTURE 25DG HERBICIDE with PURSUIT may result up to 10% reduction of grass control.

Refer to the PURSUIT label for proper weed growth stages and timing of the tankmix application. Apply this tankmix only to grasses that have not yet begun to tiller and are under optimum growing conditions. This tankmix will result in reduced grass control (particularly wild oats) if it is applied to grasses that have started to tiller and are under any kind of weather related stress conditions.

Crops treated with PURSUIT cannot be grazed or cut for hay. When tank-mixing, follow the most restrictive precautions, directions and recropping/rotation limitations that appear on both product labels.

## MIXING INSTRUCTIONS

Follow the mixing instructions below for adding VENTURE 25DG HERBICIDE to the spray tank.

1. DO NOT APPLY BY AIR. Use ground equipment only.
2. Only use sprayers in good running condition with high agitation. Ensure the sprayer is cleaned according to instructions on label of the product used prior to VENTURE 25DG HERBICIDE. Use only clean water for the spray solution. **Ensure that** all in-line **strainer** and **nozzle** screens in the sprayer are **50 mesh or coarser**. Screens of 80 mesh size or finer should not be used.
3. Begin to fill sprayer tank with clean water, and engage agitator. **Agitation must be continued throughout the entire mixing and spraying procedure.**
4. When the sprayer is three quarters full of water, add VENTURE 25DG HERBICIDE **slowly** to the spray tank. Do not enlarge the opening of the VENTURE 25DG HERBICIDE box. Remove straining screen from filler opening of spray tank before adding VENTURE 25DG HERBICIDE. (NOTE: VENTURE 25DG HERBICIDE must be added directly into the sprayer through the tank opening and **not through injector or hopper systems**, for sprayers so equipped). For shallow spray tanks where water depth is 60 cm or less, add VENTURE 25DG

---

HERBICIDE towards agitator unit and away from outlet in the bottom of the tank to enhance dispersion. If more than 1 case of VENTURE 25DG HERBICIDE is to be used, add the VENTURE 25DG HERBICIDE from all cases prior to adding other herbicides or TURBOCHARGE® Adjuvant.

5. Wait at least one minute after the last of the VENTURE 25DG HERBICIDE has been added to the tank to allow for complete dispersion of the granules. A longer agitation period may be required to disperse VENTURE 25DG HERBICIDE when using cold water from sources such as deep drilled wells.

NOTE: If MUSTER is to be tank mixed, it must be added to the spray tank before VENTURE 25DG HERBICIDE as follows: ADD MUSTER to the spray tank followed by agitation for 2 to 3 minutes. Then add the required amount of VENTURE 25DG HERBICIDE and continue agitation for at least one minute.

6. If tank mixing, add the recommended broadleaf weed herbicide next except for MUSTER.
7. Lastly, add TURBOCHARGE Adjuvant, and then continue to fill tank to desired level with water.

## ADJUVANTS

TURBOCHARGE Adjuvant must be added to the spray solution to give a final strength of 0.5% v/v (0.5 L for each 100 L of spray solution). TURBOCHARGE Adjuvant must be added to the spray tank last.

## SPRAYING INSTRUCTIONS

VENTURE 25DG HERBICIDE is not affected by rainfall if it commences two hours or more after application.

1. DO NOT APPLY BY AIR. Use ground equipment only.
2. Apply in a spray volume of 50-100 L/ha at a pressure of 275 kPa to ensure proper dispersion, spraying characteristics and performance of VENTURE 25DG HERBICIDE. Do not apply VENTURE 25DG HERBICIDE through sprayers only equipped to apply less than 100 L/ha spray solution (such as SPRAFOIL™ or SPRAY AIR™ sprayers) since mixing problems could occur.
3. When band spraying, use a proportionately smaller spray volume per hectare.
4. Flat fan nozzles of 80E or 110E are recommended for optimum coverage. Do not use floodjet nozzles or controlled droplet application equipment. Nozzles may be angled forward 45E to enhance penetration of the crop and provide better coverage. **Ensure that all in-line strainer and nozzle screens in the sprayer are 50 mesh or coarser.** Screens of 80 mesh size or finer should not be used.
5. Ensure thorough and uniform coverage of the entire leaf area of the target plants.
6. Always ensure that agitation is continued until spraying is completed, even if stopped for brief periods of time. **If the agitation is stopped for more than 5 minutes, re-suspend the spray solution by running on full agitation prior to spraying. VENTURE 25DG HERBICIDE must be sprayed within the same day of mixing.**
7. When spraying is completed and prior to using other products, ensure the sprayer tank is thoroughly rinsed with clean water to remove any remaining VENTURE 25DG HERBICIDE residues. Using a pressure washer and the addition of a detergent or AGRAL® 90 to the rinse water will enhance removal of residues.

---

® AGRAL, TURBOCHARGE and VENTURE are registered trademarks of a Syngenta Group company.  
ALLY and BLADEX are registered trademarks of E.I. DuPont de nemours & Company.  
BASAGRAN, BASAGRAN Forte, ASSIST, MERGE and POAST are registered trademarks of BASF AG, BASF Canada.  
MUSTER and PINNACLE are registered trademarks of DuPont Canada Inc.  
LONTREL amd PURSUIT are registered trademarks of Dow AgroSciences Canada Inc.  
SELECT and AMIGO are registered trademarks of Tomen Corp.  
SPRAFOIL is a trademark of D&W Industries.  
SPRAY-AIR is a trademark of SPRAY-AIR Canada Ltd.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+) )