

87.08.10

WATER SOLUBLE

WARFARIN

RODENTICIDE CONCENTRATE

COMMERCIAL

KILLS RATS AND MICE

GUARANTEE: SODIUM SALT OF WARFARIN....0.54%

READ THE LABEL BEFORE USING

KEEP OUT OF REACH OF CHILDREN

REGISTRATION NO. 19797 PEST CONTROL PRODUCTS ACT

MANUFACTURED BY:

MOTOMCO LTD.
Madison, WI 53704 U.S.A.

DIRECTIONS

HOW TO PREPARE: Dissolve 15 ml. in 1 l. of water. Pour liquid bait in glass, plastic or crockery dishes or chicken founts.

Note: Metal containers are not recommended.

HOW TO USE: Place bait in location where rats and mice feed and travel. Baiting should be maintained continuously as long as any evidence of feeding is observed. Average kill times are from 6 to 14 days, but some feeding may continue to the fourth week.

FOR NORWAY RATS(*Rattus norvegicus*) and ROOF RATS(*Rattus rattus*):
Minimum of 240 ml. of liquid should be placed in several locations, such as, along walls, in corners, and where they can

drink unobserved. Liquid station should be maintained in an uninterrupted supply for at least 10 days.

FOR HOUSE MICE(*Mus musculus*): Minimum of 240 ml. liquid station should be placed at 2-3 meter intervals. Mice travel shorter distances than rats and are more difficult to control. Liquid stations should be maintained in an uninterrupted supply for at least 15 days.

IMPORTANT

Bait placements should be examined regularly and replenished before completely empty. Should baits become sour or moldy, they must be replaced. Where infestation is likely, permanent bait station should be used and bait placed in them as often as necessary.

CAUTION

Keep away from children, pets, domestic animals and wildlife. Do not expose bait where humans, livestock, poultry, pets or wildlife can drink it.

Treated baits should be placed in locations not accessible to children, or pets, wildlife and domestic animals, or in tamper-proof bait boxes. Do not contaminate water by cleaning or equipment, or disposal of wastes. Apply this product only as specified on this label.

PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case intravenous and oral administration of Vitamin K1 combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

NOTICE TO USER: This product is to be used only in accordance with the directions on this label. It is an offense under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))