

92.01.08

BIO-SEPT

Disinfectant - Sanitizer  
Fungicide -Deodorizer

For Hospitals, Institutional, Industrial,  
School, Dairy and Other Farm Use

COMMERCIAL

GUARANTEE:

Didecyl dimethyl ammonium chloride ..... 7.5%

WARNING

POISON

IRRITANT

READ THE LABEL BEFORE USING

REGISTRATION NO. 22296  
PEST CONTROL PRODUCTS ACT

DIN 1945432

4 L          20 L          204.57 L

Manufactured by:  
CHEMPEX INDUSTRIES LTD.  
Unit #3 - 3131 Delta Ave.  
Victoria, B.C. V8Z 1A6

SANITIZING OF FOOD PROCESSING EQUIPMENT AND OTHER HARD SURFACES  
IN FOOD CONTACT LOCATIONS.

For sanitizing food processing equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces. No potable rinse water is required.

Wash and rinse all articles thoroughly, then apply a solution of 2 mL BIO-SEPT in 1 litre of water (150 ppm active). Surfaces should remain wet for at least 1 minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when used solution becomes visibly dirty. For mechanical application, used solution may not be reused for sanitizing applications.

PRECAUTIONARY STATEMENTS: KEEP OUT OF REACH OF CHILDREN.  
Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

TOXICOLOGICAL INFORMATION: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

DISPOSAL:

1. Rinse the emptied container thoroughly and add the rinsings to the treatment site.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal or reconditioning.
3. Dispose of the container in accordance with provincial requirements.
4. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environment Canada.
5. For small spills. Recover free liquid. Soak up residue with solids absorbent. DO NOT FLUSH INTO SURFACE STREAMS. INFORM ENVIRONMENT CANADA AS IN NO. 4.

DIRECTIONS FOR USE:

Apply BIO-SEPT with a cloth or mop. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the used solution becomes visibly dirty.

DISINFECTION IN HOSPITALS, NURSING HOMES, AND OTHER HEALTH CARE INSTITUTIONS: For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces, add 7 mL BIO-SEPT to 1 litre water. Apply to previously cleaned hard surfaces with a mop or cloth. At this use level BIO-SEPT is effective against *Pseudomonas aeruginosa*.

DISINFECTION IN INSTITUTIONS, INDUSTRY AND SCHOOLS: For

disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces. Add 4 mL/L of BIO-SEPT to 1 litre of water. Apply to previously cleaned hard surfaces with mop or cloth.

At 4 mL/L use level, BIO-SEPT is effective against Staphylococcus aureus, Salmonella choleraesuis and Trichophyton interdigitales, (the athlete's foot fungus).

DISINFECTION OF BARBER TOOLS: Precleaned barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 4.0 mL/L solution of BIO-SEPT.

DISINFECTION OF POULTRY EQUIPMENT, ANIMAL QUARTERS AND KENNELS: Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thoroughly cleaning by applying a solution of 4.0 mL BIO-SEPT in 1 litre of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their foods must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact food or drinking water must be rinsed with potable water before reuse.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))