

28-JAN-2005

25-FEB-2008 Notification - Change in Registrant Name

**WILSON
TURF-RITE® 2 + 2
DOUBLE STRENGTH HERBICIDE**

Contains MECOPROP and 2,4-D for use on Turf

COMMERCIAL

REGISTRATION NO. 19391 PEST CONTROL PRODUCTS ACT

GUARANTEE:

2,4-D (isomer specific) 200 g/L
Mecoprop (d-isomer) 200 g/L
(both present as amines)

**DO NOT FREEZE
SHAKE WELL BEFORE USING**

WARNING – POISON

READ THE LABEL BEFORE USING

NET CONTENTS: 4 L (9.5 L) (10 L) (20 L) (205 L)

Notification Change

~~Nu-Gro PR Inc.~~ Agrium Advanced Technologies RP Inc.
10 Craig St.,
Brantford, ON
Canada N3R 7J1

DIRECTIONS: General information: This product contains Mecoprop and 2,4-D. This combination controls a wide range of different weeds as Mecoprop is effective on certain weeds which are resistant to 2,4-D such as Black Medick, Chickweed and White Clover.

Weeds controlled: Results show in 5 to 7 days after spraying:

Annual and perennial broad-leaved weeds (bindweed*, black medick, buttercup, Canada thistle, chickweed, clover, corn spurry, curled dock, dandelion, dock, English daisy, fall hawkbit, grass-leaved stitchwort, ground-ivy (creeping Charlie), knotweed, mallow**, mossy stonecrop, narrow-leaved plantain (ribgrass), ox-eye daisy, ragweed, shepherd's purse, speedwell, thyme-leaved sandwort, wild white clover and poison ivy. This product is suitable for use on turf-grass, other established grass areas, vacant lots and along fence rows.

* Temporary Control ** Young Stages

Rate of application: Select the proper rate from this range depending on size and species of weeds present, time of year and weather conditions. DO NOT USE on Bent grass to avoid injury.

AREA	PRODUCT RATE	WATER VOLUME
100 m ²	40 - 50 mL	3 - 8 L
1 hectare	4.25 - 5.50 L	275 - 825 L

DO NOT APPLY BY AIR

Transfer System: Container larger than 20 litres: Use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

Re-entry Interval: Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treated areas until spray has thoroughly dried.

Time of application: For best results in selective weed control, the weeds should be actively growing. Therefore, application during the months of May, June and September is preferred. The effectiveness of a treatment in July and August will depend on general weather conditions. If it is hot and dry during that period, a second treatment may be necessary. If it rains within 24 hours after spraying, a repeat application may be required. Avoid mowing for several days before and after treatment. For ground-ivy (creeping Charlie), apply a spot treatment in June when the plants are young. A follow-up application in September is often necessary. For knotweed, apply in May or early June when the young plants come up. NEWLY SEEDED AREAS should not be sprayed within 4 weeks after grass emergence and then use only half the recommended rate for the first month.

Maximum application frequency to turf: For good turf/lawn management, normally two applications per year per treatment site are adequate. This does not include spot treatments.

Tank mixtures with dicamba herbicide: If a combination of this product with dicamba is required, mix 4.25 L of TURF-RITE® 2 + 2 with 275 mL of Dycleer® Liquid Herbicide (equals 100 g active dicamba) in 275 - 825 L of water for spraying one hectare. The addition of dicamba will extend the period during which prostrate knotweed can be controlled. Do not use on bent grass.

Cleaning equipment: All equipment must be cleaned thoroughly after use. Fill with hot water and detergent; flush all parts several times. Let sprayer stand with solution for 24 hours then rinse with water. It is often more convenient to use a separate sprayer with coarse nozzles for applying weedkillers.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Avoid breathing spray mist. Avoid spray drift. **Protective Clothing and Equipment: Handling the concentrate (mixing and loading):** wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes and protective eye wear (face shield or safety glasses). Rinse gloves before removal. Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5 L. **Handling the dilute spray solution (during application or repairing or cleaning equipment):** wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

Operator Use Precautions: Wear freshly laundered clothing and clean protective equipment daily. Rinse gloves before removal. Wash hands before eating, drinking, using tobacco or using the toilet. If herbicide penetrates clothing, remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry. After work, remove all clothing and shower using soap and water. Keep in original container during storage. Do not allow spray, spray mist, visible or invisible vapours or air containing mist or vapours to contact plants other than those being sprayed. If 2,4-D or MCPA susceptible plants are growing within a distance of 400 metres from the area being sprayed, spray only when there is a light breeze away from the susceptible plants. Spray drifting can be reduced by using high volumes of water sprayed under low pressure, using coarse spray nozzles.

Susceptible crops include:

beans	cauliflower	sugar beets	tobacco	turnips
cabbage	grapes	sunflowers	tomatoes	

as well as many other field, vegetable, ornamental and fruit crops. Do not contaminate any body of water, or watercourse, water used for livestock or for domestic purposes. Do not store open container near food, feedstuffs, seeds, fertilizers or pesticides including insecticides, fungicides and herbicides other than those containing 2,4-D or brush killers. Sprayers which have contained TURF-RITE® 2 + 2 should not be used to apply other pesticides to 2,4-D or MCPA susceptible crops.

FIRST AID: If on skin, wash with soap and water. **If in eyes,** flush with clean water for 15 minutes. **If irritation develops,** consult a physician or contact a Poison Control Centre immediately. **If swallowed,** drink 1-2 glasses of water to dilute the product. **DO NOT** induce vomiting. Call a doctor or Poison Control Centre immediately. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: May cause eye irritation on contact.

DISPOSAL: Triple- or pressure-rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

DYCLEER® est une marque déposée de Syngenta Crop Protection Canada Inc.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.