

2000/2005

Apache

FLY BAIT

Attracts and kills more female flies

Advanced Bait Formula contains TRICOLURE(TM) Sex Attractant
and Feeding Adjuvants

Attracts and Kills House Flies, Plus Many Other Fly Species
as Listed on this Label

Attacks Both Sexes of Flies to the Bait

Kills Quickly

CAUTION POISON

GUARANTEE:

Methomyl.....1.000%
Z-9 Tricosene.....0.025%

AGRICULTURAL

READ THE LABEL BEFORE USING

REGISTRATION NO.22303
PEST CONTROL PRODUCTS ACT

A product of

Farnam Companies, Inc.
301 W. Osborn, Phoenix, AZ
Product Code: 73009

Net Contents: 2.2 Kg

Printed in U.S.A.

9E7

PRECAUTIONS
KEEP OUT OF REACH OF CHILDREN

Hazards to humans and domestic animals: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing.

Do not contaminate feed, water, food or litter. Do not apply where poultry or animals can pick up or lick the product. Do not use in homes or where food is processed or stored. Do not feed treated garbage to livestock. Use only in areas inaccessible to food producing animals.

Environmental hazards: This bait is toxic to bees, wildlife and fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

TOXICOLOGICAL INFORMATION

Methomyl is a cholinesterase inhibitor. Poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors. Atropine is an antidote.

FIRST AID INSTRUCTIONS

In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. If swallowed, induce vomiting and get medical attention.

DIRECTIONS FOR USE

APACHE FLY BAIT is an advanced formula bait containing TRICOLURE Sex Attractant and feeding adjuvants which attract both sexes of the housefly and many other pest fly species including little houseflies, flesh flies, blow flies and eye gnats directly to the bait. Bait should be applied to specified areas of high fly population daily or as needed to reduce fly populations. Scatter bait from container or other device at the rate of approximately 250 g per 100 square meters of fly feeding area. Use APACHE FLY BAIT in and around feedlots, dairies, stables, broiler houses, hog houses, livestock barns and kennels. Bait may also be used on walkways in caged layer houses. Apply only to areas out of reach of animals.

For use in the APACHE TP BAIT STATION: Fill bait chamber to bottom of hopper by pouring 60 g of bait into hopper. Use at the rate of approximately two bait stations per 50 square meters of fly feeding area. Hang bait stations out of reach of food producing animals, or set on ground or other surface only in areas inaccessible to food producing animals, pets and children.

STORAGE AND DISPOSAL

STORAGE

Store in cool dry place inaccessible to children and pets.

PESTICIDE DISPOSAL

Follow provincial instructions for any required cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of the container in accordance with provincial requirements. For information on the disposal of unused, unwanted product and the cleanup of spills, contact the regional office of Environmental Protection, Environment Canada.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the Pest Control Products Act to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))