

13-JUL-2005 Notification received of a label change. See below for this change.

22-MAY-2007 Label correction on page 2.

07-FEB-2003

COMMERCIAL

Net Contents 20 kg

captan 7.5 DUST
Fungicide

DANGER - CORROSIVE TO EYES
POTENTIAL SKIN SENSITIZER
KEEP OUT OF REACH OF CHILDREN
AND UNAUTHORIZED PERSONNEL

READ THE LABEL BEFORE USING

CAUTION: Exposure to captan may produce long term health effects. To minimize exposure, follow directions outlined on this label.

GUARANTEE: captan 7.5% (minimum)

REGISTRATION NO. 9319 PEST CONTROL PRODUCTS ACT

Arysta LifeScience North America Inc

~~ARVESTA CORPORATION~~

100 First Street, Suite 1700

San Francisco, CA 94105

(415) 868-1103

Notification Change

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN AND UNAUTHORIZED PERSONNEL.

CORROSIVE to eyes. DO NOT get in eyes. MAY IRRITATE the skin. DO NOT get on skin. Never handle material with bare hands. Avoid spilling on skin or clothing. Do not breathe dust. Wear long sleeved and long pants, and chemical resistant gloves, goggles and NIOSH approved quarter or half mask cartridge respirator (for particulators) during mixing, loading, clean-up and repair activities. Change contaminated clothing daily and wash thoroughly before re-use. Wash hands, face and arms thoroughly after handling captan and before drinking. Shower after use or at the end of the work day. Wash clothing and gloves before re-use. Always work out of the wind during mixing and loading operations. Keep unprotected persons out of operating area or vicinity where there may be drift. Label all unplanted seed potatoes 'Poisonous to man and animals. Do not use for food or feed. These seed potatoes have been treated with Captan.'

Update

DO NOT reenter ornamentals within 4 days of treatment. DO NOT reenter all other treated areas within 48 hours of application. If re-entry into treated areas is required workers must wear long pants, long-sleeved shirt, chemical resistant gloves, work boots and goggles. No re-entry shall occur in treated greenhouses without an approved respirator unless it has been fully ventilated.

If this pest control product is to be used on a commodity that may be exported to the U.S., contact 1-866-375-4648 or www.cropro.org/.

FIRST AID

If a known exposure occurs or is suspected, immediately initiate the recommended procedure below. If further treatment is required, contact a **Poison control Centre**, a physician or the nearest hospital.

If Swallowed: Immediately give several glasses of water and induce vomiting by gagging the victim with a finger or blunt object placed on the back of the victim's tongue. Give fluids until vomitus is clear. If victim is unconscious, do not induce vomiting or give anything by mouth.

For eye contact: Immediately flush eyes with large amounts of running water for at least 15 minutes. Hold eyelids apart during flushing to ensure rinsing of the entire surface of the eye and lids with water. Get medical attention.

For Skin Contact: Flush all affected areas with plenty of water for 15 minutes. Remove contaminated clothing and shoes. Seek medical attention if skin irritation occurs. Wash clothing before re-use.

If Inhaled: Remove to fresh air. Seek medical attention if respiratory irritation occurs or if breathing becomes difficult.

EMERGENCY TELEPHONE NUMBER

All hours, 1-800-263-0984 **ONLY** for health and environmental information.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly or indirectly to water. Do not contaminate ponds, lakes streams, or rivers by cleaning of application equipment or disposal of waste.

STORAGE

Store in a cool, dry place. Protect from excessive heat. Do not contaminate food or feed.

DISPOSAL

Thoroughly empty the contents into the application device. Follow provincial instructions for any required additional cleaning of the container prior to its disposal. Make the empty container unsuitable for further use. Dispose of container in accordance with provincial requirements. For information on the disposal of unused, unwanted product contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of spills and for cleanup of spills.

DIRECTIONS FOR USE

Apply the specified amounts of this product uniformly to the foliage when air is calm, using ground equipment. Apply at the first sign of infestation and repeat 5-10 day intervals as needed to maintain control. Do not apply when weather conditions favour drift of dust from areas treated.

APPLICATION RATES

Apple: Scab, Frog-Eye Leaf Spot - 84-112 kilograms per hectare as required to maintain adequate coverage in the pre-bloom and blossoms applications. Do not apply within 7 days of harvest. Do not apply by air.

Cherry (Sour, Sweet) - Brown Rot, Leaf Spot (Shot-Hole) - 56 to 67 kilograms per hectare in the blossom and pre-harvest applications. May be applied up to 2 days before harvest of sweet cherries. May be applied up to 5 day before harvest of sour cherries. Do not apply by air.

Peach: Brown Rot and Scab - 56 to 67 kilograms per hectare in the blossom and pre-harvest applications May be applied up to 2 days before harvest. Do not apply by air.

Strawberries: Botrytis Rot - 34 to 45 kilograms per hectare beginning as soon as new growth starts. Repeat at about weekly intervals, up to harvest. Apply after each picking if conditions favour fruit rot. May be applied up to 2 days before harvest. Do not apply by air.

Cucumber: Anthracnose - Start application when runners begin to form or at the first sign of disease. Apply at a rate of 22.5 to 34 kilograms per hectare on young plants increasing to 34 to 45 kilograms per hectare on mature plants. May be applied up to 2 days before harvest. Do not apply by air.

Potato Seed Piece Treatment: Damp-Off, Seed Rot- Use at a rate of 250 grams per 25 kilograms cut seed pieces, (50 grams per 50 kilograms of cut seed pieces). Apply in a convenient container or by duster attachment over belt. Coat each piece evenly with a light film of dust. Treat pieces within 6 hours after cutting. If planting is delayed for a day or two, store treated seed pieces in open crates and provide sufficient ventilation to allow cut surfaces to dry before bagging.

Tomatoes: Anthracnose, Early and Late Blight, Gray Leaf Spot, Septoria Leaf Spot - Apply 34 to 45 kilograms per hectare at weekly intervals on young plants: 45 to 56 kilograms per hectare on mature plants. May be applied up to 2 days before harvest. Do not apply by air.

Soil and Greenhouse Bench Treatment - Apply 1.25 kilograms per 100 square metres (112 kilograms per hectare). Work into upper 7.5 to 10 centimetres of soil before planting. For Damp-off and root rot diseases of seedlings and transplants of roses, other shrubs, flowers, lawn seedbeds and vegetables such as beans, peas, celery, cabbage, eggplant, pepper and tomato. Do not apply by air.

Ornamentals: Rose-Black Spot; Carnation-Leaf Spot; Chrysanthemum-Flower Blight - Apply thoroughly prior to or at first signs of disease. Repeat application at 7 to 10 day intervals. Shorten interval during periods of frequent rain and heavy dews. Do not apply by air.

Turf (golf courses and sod farms only): Brown Patch, Damp-Off, Leaf Spot, Root Rot - Apply 1 to 1.5 kilograms per 100 square metres. Repeat at 7 to 10 day intervals as needed.

Melting-out - apply 1 to 1.5 kilograms per 100 square metres beginning when grass resumes growth in the spring (March - April). Repeat at 7 to 14 day intervals as long as cool, wet weather persists. The maximum application rate to turf (golf courses, sod farms) is 63 kgs of Captan 7.5 Dust per hectare with a maximum seasonal rate of 126.7 kgs/hectare. Do not apply to home lawns, parks, schools, sports fields and other recreational areas. Do not apply by air.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offense under the **Pest Control Product Act** to use a control product under unsafe conditions.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))