

05-APR-2005

29-JAN-2008 Notification - Change in Registrant

**LATER'S
COPPER SPRAY WETTABLE POWDER
FUNGICIDE**

Contains Fixed Copper.
Controls Anthracnose, Blight, Fireblight and Peach Leaf Curl.
For use on listed vegetable, ornamentals and fruit trees.

**READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

GUARANTIE: Copper from Copper Oxychloride 50%

DOMESTIC

250 g (200 g, 400 g)

REG. NO. 16140 P.C.P. ACT

WARNING

POISON

Notification Change

~~Spectrum Brands IP Inc.
10 Craig Street
Brantford, ON
N3R 7J1~~

Sure-Gro IP Inc.
P.O. Box 21001
Brantford ON
N3R 7W9

DIRECTIONS FOR USE:

Later's Copper Spray is concentrated and when mixed with water will provide good control for listed plant diseases. Thorough coverage of all plant parts is essential. Fill spray tank partially with water and add the required amount of **Copper Spray** while stirring. Add more water and keep agitating during filling and spraying operations. May be used up to 1 day before harvest.

3 g = 1 level teaspoon 9 g = 1 level tablespoon 1 L = approx. 1 quart

1 m = approx. 1 long pace 10 m² = 2 m x 5 m

PLANT	DISEASE
Cucumber, Melon, Pumpkin, Squash	Angular Leaf Spot, Anthracnose, Bacterial Wilt, Downy Mildew, Alternaria and Septoria Leaf Spot
Use 4 g in 1 L of water per 10 m ² . Spray at weekly intervals. For Bacterial wilt it is necessary to control Cucumber beetles.	
Potato	Early and Late Blight
Use 5 g in 1 L of water per 10 m ² , spray to ensure thorough coverage of plants. Start applications when plants are 10-20 cm high and repeat at 7-10 day intervals	
Tomato	Bacterial Canker, Early and Late Blight, Septoria Leaf Spot
Use 5 g in 1 L of water per 10 m ² , spray to ensure thorough coverage of plants. Start applications when plants are 10-13 cm high and repeat at 7-10 day intervals.	
Celery	Early and Late Blight
Use 5 g in 1 L of water per 10 m ² , spray to ensure thorough coverage of plants. Start application when plants are 5 cm high and repeat at 7-10 day intervals.	
Grape	Downy Mildew, Powdery Mildew
Mix 30 g plus 50 g of hydrated lime in 10 L of water. Apply at pre-bloom at 1.5 L of this mixture per 10 m ² and at 2 L per 10 m ² in post-bloom and cover sprays until mid-August. Do not apply on Seibel varieties.	
Sour Cherry	Brown Rot, Leaf Spot
Use 2 g plus 4 g hydrated lime per 1 L of diluted spray. Consult local spray calendar for timing of application.	
Flowering Prunus spp. Such as Cherry, Plum	Coryeum Blight, Bacterial Canker
Mix 6 g in 1 L of water. Apply before fall rains and repeat after leaves have fallen.	
Rose, Lilac, Forsythia	Bacterial Blight
Mix 6 g in 1 L of water. Apply once in October and again in January. During blight conditions in April and May, apply 2 g per 1 L of water. Repeat at 7-10 day intervals.	
Raspberry	Bacterial Blight
Use 3 g in 1 L of water to cover 10 m ² . Apply once before fall rains and 3 times using 2 g per 10 L of water to cover 10 m ² beginning with budburst and at 14-28 day successive intervals. Do not apply within one day of harvest.	
Onion	Downy Mildew
Use 3 g per 1 L of water to cover 10 m ² . Apply 10-14 days throughout the growing season, beginning in mid-May for over-wintered crops and mid-June for spring seeded crops. Do not apply within one day of harvest.	
Peach, Apricot (B.C. only)	Coryeum Blight, Peach Leaf Curl
Apply 9 g per 1 L of water in September AFTER the harvest is completed.	
Pear (B.C. only)	Fireblight (see below)

To reduce blossom infection, apply 3 g per 1 L of water when blossoms open. In case of hail damage immediately repeat this treatment. Do not apply on Anjou.	
Cotoneaster, Crabapple, Hawthorne, Mountain Ash, Quince	Fireblight (see below)
Apply 1.5 g in 1 L of water several times during blossoming; also repeat during summer when bacteria spread rapidly in warm rainy weather	
Arborvitae, Cedar, Fir, Juniper, Pine, Spruce	Leafblight, Needlecast, Tipblight, Twigblight
Apply 4 g in 1 L of water when new growth starts in spring and repeat 3 to 4 times at 10 day intervals. Further sprays are advisable during prolonged rainy periods.	
FIREBLIGHT may be spread to healthy plants by insects such as aphids and leafhoppers. Diseased twigs which appear to be 'scorched' must be cut off 15 cm below the infected part and burned; disinfect pruning shears after each cut in solution of 1 part bleach to 4 parts of water.	

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. After handling, wash thoroughly with soap and water before eating, drinking or smoking and change to clean clothing. Avoid contamination of feed or food.

ENVIRONMENTAL HAZARDS: Do not apply or allow drift to areas occupied by animals, or to streams, lakes or ponds to protect wildlife.

FIRST AID:

IF SWALLOWED, give two glasses of water to drink and induce vomiting by inserting finger into mouth and touching back of the throat. Repeat until vomit fluid is clear. Contact a doctor or poison control centre IMMEDIATELY. Never give anything by mouth to an unconscious person.

IF IN EYES, flush with water for 15 minutes and get medical attention.

IF ON SKIN, remove contaminated clothing and wash skin with soap and water or contact poison control centre IMMEDIATELY.

Take container, label or product name and P.C.P. Reg. No. with you when seeking medical attention.

STORAGE: Keep product in original container. Store in a cool, dry place away from food.

DISPOSAL: Do not reuse empty container. Wrap container and discard in household garbage.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

NOTICE TO BUYER: Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.