MATAVEN (R) L

AGRICULTURAL

HERBICIDE

Emulsifiable concentrate for control of Wild Oats

Net Contents: 10 L

GUARANTEE: flamprop-M-methyl......52.5 g/L

REGISTRATION NO: 18799
PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING

WARNING

POISON IRRITANT

DU PONT CANADA

Regency Centre #105 333 - 25th Street E. Saskatoon, Saskatchewan S7K 0L4

Box 2300, Streetsville Mississauga, Ontario L5M 2J4

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID:

This product contains petroleum distillate. If ingested do not induce vomiting. If patient is unconscious, give him air. Keep patient prone and quiet. Transport the patient immediately to the nearest physician.

Skin Contact: If spilled on the skin, remove all contaminated clothing at once. Thoroughly wash skin with soap and water. Contact physician. Wash contaminated clothing before reuse.

For Eyes: If the material gets into the eyes, flush immediately with running water for at least 15 minutes. Contact physician.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. Product is harmful if swallowed. Avoid contact of this product with eyes. When handling, the use of safety goggles is recommended. Avoid inhalation of vapors. Keep product in original closed container and store away from food and feed. Wash hands and exposed skin before eating, drinking, smoking and after work. Flush concentrate from skin and eyes with water immediately after contact.

DISPOSAL AND DECONTAMINATION:

- 1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
- 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- 3. Make the empty container unsuitable for further use.
- 4. Dispose of the container in accordance with provincial requirements.
- 5. For information on the disposal of unused, unwanted product, and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

PAMPHLET

MATAVEN (R) L

AGRICULTURAL

HERBICIDE

Emulsifiable concentrate for control of Wild Oats

GUARANTEE: flamprop-M-methyl......52.5 g/L

REGISTRATION NO: 18799 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED PAMPHLET BEFORE USING

WARNING

POISON

IRRITANT

DU PONT CANADA
BOX 2300, STREETSVILLE
MISSISSAUGA, ONTARIO
L5M 2J4
1-800-667-3925

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the PEST CONTROL PRODUCTS ACT to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID:

This product contains petroleum distillate. If ingested do not induce vomiting. If patient is unconscious, give him air. Keep patient prone and quiet. Transport the patient immediately to the nearest physician.

Skin Contact: If spilled on the skin, remove all contaminated clothing at once. Thoroughly wash skin with soap and water. Contact physician. Wash contaminated clothing before reuse.

For Eyes: If the material gets into the eyes, flush immediately with running water for at least 15 minutes. Contact physician.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. Product is harmful if swallowed. Avoid contact of this product with eyes. When handling, the use of safety goggles is recommended. Avoid inhalation of vapors. Keep product in original closed container and store away from food and feed. Wash hands and exposed skin before eating, drinking, smoking and after work. Flush concentrate from skin and eyes with water immediately after contact.

DISPOSAL AND DECONTAMINATION:

- 1. Rinse the emptied container thoroughly and add the rinsings to the spray mixture in the tank.
- 2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
- 3. Make the empty container unsuitable for further use.
- 4. Dispose of the container in accordance with provincial requirements.
- 5. For information on the disposal of unused, unwanted product, and the cleanup of spills, contact the regional office of Conservation and Protection, Environment Canada.

CROP INFORMATION:

MATAVEN L is a post-emergent herbicide for the control of wild oats in: wheat (provincially recommended varieties of spring, durum, winter wheat, triticale, canary grass, sunflowers and certain forage crops (seed production only).

TIMING OF APPLICATION:

MATAVEN L can be applied to wild oats from the 3-leaf stage to the 6-leaf stage. For optimum control and maximum yield response, apply Mataven L when the majority of the wild oats have 3-4 leaves.

TIME AND RATE OF APPLICATION:

| CROP | CROP STAGE | LITRES MATAVEN L PER HECTARE | REMARKS |
|---|--|---------------------------------|---|
| Wheat (provincially recommended varieties of spring, durum, Winter), Canary grass Triticale | Not beyond 6 - leaf stage | 5.0 | |
| Sunflowers | Not beyond 6-leaf stage but before dense crop canopy forms | 5.0 - 6.5 | Use the higher rate when wild oat pressure is heavy and for band application over crop row. Band application should be combined with interow cultivation. |

| CROP | CROP STAGE | LITRES MATAVEN L PER HECTARE | REMARKS |
|--|--------------------|---------------------------------|--|
| Alfalfa, Red Clover, Birds-foot trefoil Sainfoin, Cicer milk vetch | In year of seeding | 5.0 | Do not harvest or graze forages in year of treatment |
| Bromegrass, Meadow Fescue, Creeping Red Fescue, Russian Wild Ryegrass, Intermediate Wheatgrass, Crested Wheatgrass | In year of seeding | 5.0 - 7.5 | Use the high rate when grown without a companion crop. Use the low rate when grown with a companion crop for which Mataven L is registered. Do not harvest for forage or graze in year of treatment. |

MATAVEN L PLUS GLEAN ** TANK MIX (WHEAT ONLY):

Apply Mataven L at 5 litres per hectare tank mixed with Glean at 15 to 30 grams per hectare for the control of wild oats and broadleaf weeds in spring and durum wheat. For optimum control, apply this tank mixture when the wild oats are in the 3-4 leaf stage (majority in the 4-leaf stage).

DO NOT ADD SURFACTANTS OR OTHER ADDITIVES TO THIS TANK MIX. CONSULT THE GLEAN LABEL FOR DETAILED INFORMATION AND SPECIAL RESTRICTIONS.

MIXING INSTRUCTIONS:

- 1. Fill the sprayer tank 1/2 full with water. Engage gentle agitation.
- 2. Add Glean (if applicable) and ensure it is completely suspended in the water.
- 3. Add Mataven L. Be certain Glean is fully mixed before adding Mataven L.
- 4. Finish filling spray tank and maintain gentle agitation during mixing and spraying.
- 5. Do not let contents stand without agitation system engaged.

APPLICATION INSTRUCTION:

Ground Application:

Add Mataven L to not less than 100 litres of water per hectare, using flat fan spray tips at a minimum pressure of 300 kPa. Use 50 mesh or larger in-

line and nozzle screens. Directing the spray forward 45 Deg. contributes to better coverage. If crop or weed canopy is dense, increase water volume to improve coverage and weed control.

Aerial Application:

Add Mataven L to not less than 15 litres of water per hectare. Do not spray in winds greater than 15 km/hr. Do not allow spray drift to contaminate other crops. Application should not be made by air when oat crops are adjacent and downwind of the treated area. Do not apply Mataven L plus Glean by air.

USE PRECAUTIONS:

- . Do not tank mix with any herbicide other than Glean.
- . Do not tank mix with spray adjuvants, surfactants or liquid fertilizer.
- . Allow 4 days between an MCPA or bromoxynil application and an application of Mataven L.
- . Allow 7 days between a 2,4-D or dicamba application and an application of Mataven L.
- . Rainfall within 2 hours of application may reduce effectiveness of Mataven L.
- . Protect Mataven L from freezing.
- * Registered trademark of Shell Canada Limited DuPont Canada Inc. is the registered user.
- ** Glean is a registered trademark of E.I. DuPont de Nemours and Co.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.

+))